

**At-home use of app-based mindfulness for children: A randomized active-controlled trial**

Isaac N. Treves<sup>a,b</sup>, Halie A. Olson<sup>a</sup>, Ola Ozernov-Palchik<sup>a</sup>, Cindy E. Li<sup>b</sup>, Kimberly L. Wang<sup>a</sup>,  
Xochitl M. Arechiga<sup>a</sup>, Simon B. Goldberg<sup>c,d</sup>, and John D.E. Gabrieli<sup>a</sup>

<sup>a</sup>Department of Brain and Cognitive Sciences and McGovern Institute for Brain Research,  
Massachusetts Institute of Technology, 43 Vassar Street, Cambridge, MA 02139, USA

<sup>b</sup>Hock E. Tan and K. Lisa Yang Center for Autism Research at Massachusetts Institute of  
Technology, Cambridge, MA

<sup>c</sup>Department of Counseling Psychology, University of Wisconsin-Madison, Madison, WI, USA

<sup>d</sup>Center for Healthy Minds, University of Wisconsin-Madison, Madison, WI, USA

**Author Note**

IN Treves <https://orcid.org/0000-0001-7802-6145>

Correspondence should be addressed to Isaac Treves, 43 Vassar Street 46-4077,  
Cambridge, MA 02139, 1-608-770-6805, [treves@mit.edu](mailto:treves@mit.edu)

**Declarations****Compliance with Ethical Standards**

This project received Institutional Review Board approval at the Massachusetts Institute of  
Technology (MIT).

**Consent**

We obtained informed consent from the parents of all participants as well as assent from the  
children before they participated in the study.

**Conflicts Of Interest**

The authors have no relevant financial or non-financial interests to disclose.

**Funding Statement**

This research was funded by the Chan Zuckerberg Initiative as part of the Reach Every Reader Project, National Institutes of Health (F32-HD100064 to OO), and the National Science Foundation Graduate Research Fellowship (Grant No. 1745302 to HO). SBG was supported by the National Center for Complementary & Integrative Health of the National Institutes of Health under Award Number K23AT010879.

**Data, Materials and Code Availability**

Data and code will be made publicly available on publication at <https://osf.io/23vax>.

**Author Contribution**

HAO, OO, INT, JDG contributed to the study conception and design. Material preparation and data collection were performed by CEL, XA, KW, HAO, OO. Analysis was performed by INT, with consultation by SBG. The first draft of the manuscript was written by INT and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

**Author CRediT**

Isaac Treves: Conceptualization, analysis, writing-original draft preparation, supervision. Halie Olson: Conceptualization, project administration, funding acquisition, writing-review and editing. Ola Ozernov-Palchik: Conceptualization, project administration, funding acquisition, supervision. Cindy Li: data curation, writing-original draft preparation, investigation, methodology. Kimberly Wang: project administration, writing-review and editing, investigation.

Xochitl Arechiga: project administration, investigation. Simon Goldberg: analysis, writing-review and editing. John Gabrieli: supervision, resources, writing-review and editing.

### **Acknowledgements**

We thank our testers: Amanda Miller, David Bates, Ross Weissman, Joohee Baik, June Okada, William Oliver, and Harriet Richards. We also thank Yesi Camacho Torres, Hope Kentala, Natalie Gardino, and Jeff Dieffenbach for helping to run the study. We thank the Los Angeles Unified School District (LAUSD) Achievement Network for help with recruiting families. We thank Learning Ally for help with recruitment and for providing the audiobook interventions. Lastly, we thank Inner Explorer for providing the intervention.

**Recommended citation:** Treves, I. S., Olson, H. A., Ozernov-Palchik, O., Li, C. E., Wang, K. L., Arechiga, X. M., Goldberg, S. B., & Gabrieli, J. D. E. (in press). At-home use of app-based mindfulness training in children: A randomized active-controlled trial. *Mindfulness*.

### Abstract

**Objectives:** School-based mindfulness interventions in children have shown benefits to child well-being. Here, we investigated the effectiveness of a remote, app-based mindfulness intervention for promoting well-being in children.

**Methods:** We conducted a randomized controlled trial (RCT) with two control groups to examine the effects of an 8-week mindfulness intervention in U.S. children ages 8-10. We compared pre-post effects between a mindfulness intervention using the *Inner Explorer* app, and two audiobook control interventions. The 279 children who participated in the interventions were assessed on self-report measures of anxiety and depression symptoms, perceived stress and trait mindfulness and we also collected parental reports.

**Results:** Over 80% of children completed the intervention in each condition. There was evidence for reduced self-perceived stress in children and reduced negative affect in children by parental reports using the mindfulness app, but no significant reduction for anxiety or depression symptoms. In general, between-group effect sizes were small ( $d_s < 0.45$ ). Regular use, defined as at least 30 days of mindfulness practice within the study period, was associated with reduced child negative affect by parental reports, as well as reduced parental stress and child self-perceived stress.

**Conclusions:** These findings suggest that home use of a mindfulness app in young children can have a positive impact on children's emotional well-being if the app is used regularly, specifically for at least 30 days in the 8-week study period. Strategies aimed at promoting regular use of the mindfulness app at home could lead to even better outcomes for children.

**Preregistration:** Preregistered on OSF at <https://osf.io/23vax>

*Keywords:* Children, mindfulness, stress, mental health, app, RCT

Mindfulness is often defined as an accepting, open-minded attention to the present moment (Creswell, 2017). Mindfulness interventions in children typically consist of in-person trainings in mindfulness practices such as breath awareness, body scans, and journaling (Kabat-Zinn, 2003). These practices are intended to help children be fully aware of their experiences and to adopt an open and accepting stance toward them. Although mindfulness practices vary, they share a focus on training attention, building emotion regulation skills to effectively manage stress, and gaining self-knowledge (Greenberg & Harris, 2012). Further, some practices aim to build empathy and compassion (Kabat-Zinn, 2005). Mindfulness interventions in children have shown a host of benefits, including reduced psychopathology (Zoogman et al., 2015), reduced stress and negative feelings (Bauer et al., 2019), increased prosociality (Flook et al., 2015), improved attention (Bauer et al., 2020), and improved academic outcomes (Bakosh et al., 2016). A meta-analysis of randomized controlled trials (RCTs) in children and adolescents found decreases in anxiety relative to active control groups (Dunning et al., 2022). Due to these benefits, mindfulness has increasingly been used in schools to prevent and reduce stress and stress-related mental health and behavioral problems (e.g., mindfulness-based social emotional learning (MBSEL)) (Greenland, 2010).

Mindfulness may be beneficial for children because it teaches self-regulatory skills, which have been linked with both readiness to learn (Diamond & Lee, 2011; Mrnjaus & Krmeta, 2014; Schonfeld et al., 2015) and decreased childhood stress (Lupien et al., 2009; Miller et al., 2011; Repetti et al., 2002). Two key types of self-regulatory skills are emotion regulation, the ability to monitor, evaluate, and modify emotional reactions (Thompson, 1994) and attention regulation, the ability to flexibly adjust attention (Posner, 2011). Emotion regulation in children is associated with better academic performance, higher social competence, fewer behavior issues

and more (Beauchaine et al., 2007; Graziano et al., 2007; Spinrad et al., 2006). Attention regulation is related to lower instances of childhood ADHD, mood disorders, anxiety (e.g. Emerson et al., 2005), greater interpersonal success (Rotenberg et al., 2008), and future self-regulatory abilities and outcomes (Friedman et al., 2007; Miller et al., 2011). Despite the positive long-term impacts of self-regulation, children are rarely explicitly taught how to pay attention and regulate their emotions (Semple et al., 2017). Critically, mindfulness trainings in children teach these regulatory skills “from the inside out” by focusing on mental experiences, such as thoughts, emotional states, the breath, and other bodily sensations (Semple et al., 2010). Mindfulness interventions also incorporate attentional training, e.g., repeated gentle redirection of attention to the breath (Lutz et al., 2008). In summary, mindfulness training may lead to the development of healthy self-regulation in children, resulting in improvements in academic outcomes (Bakosh et al., 2016), social outcomes (Flook et al., 2015), and mental health outcomes (Dunning et al., 2022).

Mindfulness instruction for children has typically been delivered in schools and other settings with an adult instructor controlling the content, pace, and duration of instruction. Another potential approach is to use remote, app-based mindfulness programs with children, which could allow for more widespread mindfulness instruction that reaches many diverse children for a low cost. App-based interventions may be particularly useful for reaching children and adolescents who are increasingly involved with the digital world. The possibility of such a highly scalable and wide-reaching intervention is especially salient given evidence for increasing risk for mental health difficulties in children in the USA (Centers for Disease Control and Prevention [CDC], 2022). A recent survey found that 95% of adolescents own smartphones (Anderson & Jiang, 2018) and they spend over two-and-a-half hours per day on their phones

(Rideout, 2015). Some children and adolescents already seek mental health support on their phones (Rideout & Fox, 2018). Indeed, a review of 71 digital health interventions in children and adolescents found good retention (Liverpool et al., 2020), and another review identified improvements in mental health outcomes (e.g. depression and anxiety) (Grist et al., 2019) (although many of these interventions are internet-based CBT).

Beyond feasibility, an important open question is whether app-based mindfulness interventions are effective for children. Mindfulness apps usually deliver content in short (e.g., ten-minute) recordings, contrasting with in-person workshops where sessions may be longer. Despite this difference, app-based interventions have been efficacious in adults. App-based interventions have been shown to decrease distress (Goldberg et al., 2020; Hirshberg et al., 2022), improve trait mindfulness (Flett et al., 2019), and improve various facets of well-being including mood (Economides et al., 2018) and positive affect (Mahlo & Windsor, 2021). A meta-analysis of 66 smartphone intervention studies found significant decreases in stress and in anxiety and depression symptoms (Linardon et al., 2019).

To date, no quantitative studies have been conducted on the use of app-based mindfulness interventions in pre-adolescent children (see Nunes et al., 2020; Puzia et al., 2020; Tunney et al., 2017 for qualitative reports). However, several studies have been conducted with adolescents, with positive results. A pilot study of a six-week mindfulness app intervention in a normative population of adolescents found that adolescents who used the app reported healthier weight-related behaviors (Turner & Hingle, 2017). A larger study with 80 adolescents found that the app intervention reduced rumination, with effects persisting for up to 12 weeks after the intervention (Hilt & Swords, 2021). Finally, a three-week randomized controlled trial (RCT) with 152 adolescents (with a control mood monitoring intervention) found that the mindfulness

intervention resulted in greater reductions in rumination compared to the control condition (Hilt et al., 2023). This research suggests that app-based mindfulness interventions are effective for adolescents, and raises the question as to whether and under what circumstances such interventions are effective for pre-teen children.

While all studies so far in children have tested the efficacy of mindfulness interventions using trained adult instructors, in practice, many schools choose to implement mindfulness instructions using app-based programs. One such program, *Inner Explorer*, consists of mindfulness practices recorded by a diversity of speakers, and was developed for multiple age ranges including 8-10 year olds. *Inner Explorer* has been implemented in over 3,000 schools with 2 million students across 50 states (Inner Explorer, 2023). In one study, 383 1<sup>st</sup> – 5<sup>th</sup> grade students were randomized by classroom to either Inner Explorer, consisting of 10 weeks of daily 10 minute recordings, or waitlist control (Bakosh, 2013). Teachers endorsed high adherence (completing 95% sessions) and no disruption to daily routines. Students showed improvements in math scores as well as grade point average when compared to the waitlist control. Further analysis of these data showed particular academic improvements in a subgroup of students with disabilities (Dunlap, 2022). A non-randomized study in 191 3<sup>rd</sup> graders also showed academic improvements, and, additionally, decreases in behavioral disruptions (Bakosh et al., 2016). However, two studies without control groups found no pre-post changes in academic, mental health, or behavioral outcomes (Goodman, 2019; Strickland, 2023). Thus far, there have been no randomized active-controlled trials of *Inner Explorer*, and it has only been implemented in school settings. In general, there is sufficient feasibility and acceptability evidence to support the use of *Inner Explorer* with children, and the hands-off audio recordings may be suitable for applications outside the classroom.

In the present study, we sought to rigorously test whether *Inner Explorer* delivered remotely is beneficial for children, and thus shed light on the effectiveness of app-based mindfulness interventions for children generally. In addition, one of our motivations for remote administration was the COVID-19 pandemic and associated restrictions. From July 2020 to January 2022, we administered a remote mindfulness intervention using the *Inner Explorer* app (Inner Explorer, 2023) to children from ages 8-10 years. Children were recruited to participate in a remote at-home audiobook program aimed at enhancing reading skills and then randomized to one of three conditions: (1) the audiobook program (Audiobook-only condition); (2) the audiobook program with additional personalized support (Audiobook-scaffolded condition); or (3) the mindfulness program (Mindfulness condition) (Ozernov-Palchik et al., 2022). We followed recommendations for a rigorous RCT design (Davidson & Kaszniak, 2015) including an active control condition, intent-to-treat analyses, and preregistration of hypotheses. Our main hypothesis was that the children in the mindfulness intervention would experience greater decreases in stress and anxiety than children in the audiobook interventions. We also assessed a range of other outcomes including negative affect, which may be reduced in children with greater trait mindfulness (Treves et al., 2023) and has been reduced in school mindfulness programs for children (Vickery & Dorjee, 2016).

Of note, this study was designed to evaluate the effects of remote audiobook interventions, and mindfulness was included as an active control condition. Mindfulness may be a suitable control for an audiobook intervention, as both interventions involve listening to recordings over a period of multiple weeks, but the mindfulness intervention was not hypothesized to influence reading skills. We believe this comparison was equally apt for studying whether changes in mental health outcomes like stress and anxiety were specific to the

mindfulness intervention as the reading-related interventions were not hypothesized to enhance mental health outcomes. For context, the interventions took place during the height of the COVID-19 pandemic in the US, when social and academic restrictions were in place.

## Method

### Participants

We analyzed the pre-intervention (baseline) and post-intervention data from children and their parents in a remote audiobook intervention study, collected from July 2020 through January 2022. Third and fourth-grade children across the United States were recruited through Facebook ads, flyers, school partnerships, and through word of mouth. We aimed to recruit children from lower socioeconomic status (SES) backgrounds and therefore reached out to school districts with high percentages of students eligible for free/reduced lunch and targeted Facebook ads to lower income zip codes across the country (for more on recruitment strategies see Ozernov-Palchik et al., 2022). As a number of schools we contacted had significant numbers of Spanish-speaking families, we provided parents with the opportunity to indicate whether Spanish was their preferred language for communication. The research team included bilingual staff, fluent in Spanish, who translated all written parent communication and assessments into Spanish, and were available to communicate in Spanish with parents who indicated that was their preferred language.

A total of 1020 participants were assessed for eligibility, and 314 were randomized into the three conditions of the study: Mindfulness ( $n = 101$ ), Audiobook-only ( $n = 105$ ), and Audiobook-scaffolded ( $n = 108$ ) conditions (Fig. 1). Of the 314 participants randomized, 279 completed pre-test and were given the interventions. We collected demographic information

regarding child gender, age, grade, mental health diagnosis, parent education level and annual household income. Parents reported their child's gender, and we did not inquire about sex assigned at birth. Therefore, we utilized gender to obtain scores for the self-report measures that were normed by sex.

----- Fig. 1 -----

The 279 children (157 male) with demographic information were on average 9 years, 5 months old ( $SD = 6$  months, range 8 years, 0 months to 10 years, 5 months). The majority of children came from the states of Georgia, Massachusetts, California, and Texas. The median income of our obtained sample was 80,000-120,000 US Dollars. Median income in the United States in 2021 was 70,784 US Dollars (*census.gov*). Demographics by group are shown in Table 1.

----- Table 1 -----

### **Procedure**

The entirety of the study was conducted remotely. Potentially interested parents filled out an eligibility screener and were invited to participate in a pretest session (pre-intervention) if initial inclusionary criteria were met (child speaks English proficiently, has normal or corrected-normal hearing, has access to a computer or tablet at home, and has a parent that speaks English or Spanish). Further inclusionary criteria required that the child achieve a standard score of 80 or above on the Kaufman Brief Intelligence Test (KBIT) Matrices subtest, a standardized nonverbal IQ assessment (Kaufman, 2004).

Eligible participants whose parents opted-in to the study were block randomized into one of three experimental conditions, each lasting eight weeks. We assigned groups based on a list of block-randomized integers 1-3, sampled from a normal distribution.

We obtained informed consent from the parents of all participants as well as assent from the children over Zoom before they participated in the study.

### ***Interventions***

The mindfulness intervention consisted of an 8-week Mindfulness Based Stress Reduction (MBSR)-inspired curriculum, delivered remotely through the *Inner Explorer* smartphone app (Inner Explorer, 2023). *Inner Explorer* customized their 90-day program to 40 days for the purposes of best comparison to the 8-week audiobook interventions (as well as MBSR and the majority of mindfulness interventions). Parents of children assigned to this condition were instructed to encourage their children to engage in the practices as laid out in the *Inner Explorer* app. The program entailed approximately 10 minutes of practice per day, five times a week, over the course of the 8-week-long intervention phase. Types of practices include focused attention to breath, body scans, and gratitude and loving-kindness practices (also called ‘heart-centered’ practices by *Inner Explorer*). Support from research staff within the Mindfulness condition was minimal and was limited to video materials about the study, the app, and mindfulness in general, technical help if there were problems downloading or using the app, and weekly one-page digests briefly summarizing the past week’s practices and recommending an optional mindfulness exercise for the weekend. Parents were instructed to have their children follow the 40 practices in order and were advised to ‘catch up’ on the week’s practices over the weekend if needed. All participants were given continued access to the app at the end of the study.

There were two reading interventions, Audiobook-only and Audiobook-scaffolded. Children in the Audiobook-only intervention received unlimited access to audiobooks via the *Learning Ally* web-based platform (Learning Ally, 2023), curated based on their listening comprehension level. Children in the Audiobook-scaffolded intervention also received audiobooks and recommendations, as well as one-on-one 30-minute online sessions with a learning facilitator twice per week, focused on improving their listening comprehension strategies and supporting their intervention adherence.

For the mindfulness intervention, we recorded number of days practiced from the *Inner Explorer* app. Participants were assigned one ten-minute practice per weekday for eight weeks, so perfect adherence would be 40 days practiced. We also collected parental reports of number of practices completed with a categorical, ordinal question. For the reading intervention, we collected listening minutes from the *Learning Ally* app.

## **Measures**

We obtained child self-reports via questionnaires administered over Zoom, a secure video chat platform, during testing sessions that involved a battery of language and cognitive assessments (Ozernov-Palchik et al., 2022). Experimenters read the assessment questions to the children who followed along on their computer screen and answered aloud. Testing sessions typically ranged between 1-2 hours, and there were 2-3 pretest sessions, as well as 1-3 post-test sessions, over the course of the study. Participants found out their group assignment before pre-testing. Testers were blind to group assignment during pretest and during post-test up until the final questionnaire (which was only administered to children in the reading interventions). When possible, the same tester was assigned to a child for multiple sessions. We also obtained parent-reports of child behavior through online questionnaires administered remotely through Research

Electronic Data Capture (REDCap). REDCap is a secure, web-based application designed to support data capture for research studies (Harris et al., 2009). Parents were compensated \$5.00 for every questionnaire completed (10 total questionnaires possible over the course of the study). Children were compensated \$20.00 per hour of testing. Questionnaires were available in Spanish and English for parents and in English for children.

### *Child Measures*

**Perceived Stress.** To measure stress, we administered the Perceived Stress Scale for Children (PSS-C) (White, 2014). This self-report measure consists of 13 items on a 4-point Likert scale ranging from ‘never’ to a ‘lot.’ The questions assess perceived stress related to time pressure, academic performance, and relationships with family and friends. A higher score indicates a greater level of perceived stress. For this study the Cronbach’s Alpha was 0.60, and the McDonald’s Omega was 0.52. Perceived stress was one of the two primary outcomes in our preregistration.

**Depression & Anxiety.** To measure child-reported anxiety and depressive symptoms we used the 25-item Revised Child Anxiety and Depression Scale (RCADS-25-C) (Ebesutani et al., 2012), which contains the following scales: Anxiety Total scale, Depression Total scale, Total scale. We omitted the Total scale from analysis per our preregistration. Items are based on Diagnostic and Statistical Manual of Mental Disorders–Fourth edition (DSM-IV) criteria (American Psychiatric Association [APA], 1994). Items in the Anxiety Total scale measure a “broad anxiety” dimension, assessing a variety of anxiety symptoms. Items in the Depression Total scale measure symptoms of major depressive disorder. Higher scores represent a greater degree of symptoms. All items are rated on a 4-point Likert scale ranging from ‘never’ to ‘always.’ Raw summative scores from each of the scales are converted into *T*-scores normed by

age and sex, with the following ranges: low severity (0-64), medium severity (65-70), and high severity (>70). *T*-scores of medium severity are considered to be “borderline clinical threshold” whereas *T*-scores of high severity are considered to be “above clinical threshold” (Chorpita et al., 2000). For this study the Cronbach’s Alpha values were 0.78 and 0.69 for the Anxiety Total and Depression Total scales respectively, and the McDonald’s Omega values were 0.81 and 0.72, respectively. The Anxiety Total scale was one of the two primary outcomes in our preregistration.

**Negative Affect.** To measure children’s affect we administered a brief 13-item questionnaire (Panorama Education, 2015). Children rated the degree to which they felt each of thirteen emotions in the past week, using a 5-point Likert scale ranging from ‘almost never’ to ‘almost always.’ We calculated a Negative Affect score from the seven items asking about negative affect: mad, bored, lonely, sad, nervous, worried, and afraid. This Negative Affect factor was supported by confirmatory factor analyses (see Supplement). The composite score ranges from 7 to 35, with higher scores representing more negative affect. The Cronbach’s Alpha for this scale was 0.71, the McDonald’s Omega was 0.78.

**Mindfulness.** To assess child trait mindfulness we administered the Child and Adolescent Mindfulness Measure (CAMM) (Greco et al., 2011). This self-report scale consists of 10 items querying the frequency of non-mindful thoughts or behaviors on a 5-point Likert scale from ‘never true’ to ‘always true.’ All items are negatively worded and reverse-scored. Higher scores represent greater levels of mindfulness. Specifically, the authors describe the scale as measuring both awareness of the present and the degree to which one has a nonjudgmental attitude towards one’s thoughts and feelings, which includes not suppressing or avoiding them. In this study the Cronbach’s Alpha was 0.74, the McDonald’s Omega was 0.78.

### *Parent Measures*

**Negative Affect.** To measure child affect we asked parents to rate the degree to which their child felt worried, frustrated, stressed, sad, mad, calm, and happy, using a 4-point Likert scale, ranging from 'not at all' to 'very.' We reverse-scored calm and happy and calculated a composite Negative Affect score ranging from 0 to 21, with higher scores signifying more negative affect and less positive affect. The Cronbach's Alpha for this scale was 0.82, and the McDonald's Omega was 0.88.

**Prosociality.** To assess child prosociality we asked parents to rate how often their child is helpful when asked, how often their child is helpful without being asked, and how often their child has done something kind. Parents answered each question on a 4-point Likert scale ranging from 'not at all' to 'very.' Scores ranged from 0 to 9, with higher scores indicating more prosociality. We calculated a prosociality composite by summing the raw scores from the three items. The Cronbach's Alpha for this scale was 0.80, and the McDonald's Omega was 0.81.

**Perceived Stress.** In order to measure parents' own perceived stress parents completed the ten-item adult self-report Perceived Stress Scale (PSS) (Cohen et al., 1983; Cohen, 1988) which assesses their overall perceived stress over the past month. Items are rated on a 5-point Likert scale from 'never' to 'very often,' with four items that are reverse-scored. Higher scores indicate greater levels of perceived stress. In this study internal reliability was demonstrated with an Cronbach's Alpha of 0.87, and the McDonald's Omega was 0.91.

**Depression & Anxiety.** In addition to the child's self-report of their anxiety and depressive symptoms, we also obtained a parent report of their child's anxiety and depressive symptoms, assessed with the 25-item Revised Child and Anxiety Depression Scale, Parent Form (RCADS-25-P) (Chorpita et al., 2000; Ebesutani et al., 2017). This scale consists of the same

scales and items that comprise the RCADS-25-C: Anxiety Total and Depression Total scales (see above). The only difference is that the items are worded to ask the parent about their child's behavior. As is the case with the child scale, items are based on DSM-IV (APA, 1994). Higher scores represent a greater degree of symptoms. All of the items are rated on a 4-point Likert scale ranging from 'never' to 'always.' Raw summative scores from each of the scales are converted into *T*-scores normed by child age and sex, with the following ranges: low severity (0-64), medium severity (65-70), and high severity (>70). *T*-scores of medium severity are considered to be "borderline clinical threshold" whereas *T*-scores of high severity are considered to be "above clinical threshold" (Chorpita et al., 2000). For this study the Cronbach's Alphas were 0.75 and 0.77 for the Anxiety Total and Depression Total scales respectively, and the McDonald's Omegas were 0.80 and 0.81, respectively.

**Executive Functioning.** We administered an 86-item parent-report of the child's executive functioning, the Behavior Rating Inventory of Executive Function-Parent Form (BRIEF) (Gioia et al., 2000). Although the BRIEF has several subscales and composite indices, we chose to use the Global Executive Composite (GEC) as a summary measure of executive function, as well as the Behavioral Regulation Index (BRI). Higher scores on the GEC indicate a greater degree of difficulty with executive functioning (e.g., needs help from an adult to stay on task). Higher scores on the BRI indicate a greater degree of difficulty with behavioral regulation (e.g., overreacts to small problems). Raw scores are summed and transformed into *T*-scores normed by age and sex. *T*-scores  $\geq 65$  are considered clinically significant (Gioia et al., 2000). In this study the Cronbach's Alpha was 0.97 for the GEC, and 0.94 for the BRI, and the McDonald's Omegas were 0.98 and 0.96, respectively.

## Data Analyses

Data were analyzed using intention-to-treat principles (i.e., participants were not excluded based on engagement) (Polit & Gillespie, 2010). We assessed differences in demographics and outcomes by group at baseline using ANOVAs, and in the case of significant differences, we used Tukey tests for isolating the group differences driving the effect. We assessed differences in completion by group using logistic regression. We assessed whether demographics or baseline outcome measures predicted completion using logistic regression. Last, we assessed correlations between outcome variables at baseline (Supplementary Table S1).

We used ANOVAs to test for group by time effects, comparing Mindfulness vs Audiobook-only groups, and Mindfulness vs Audiobook-scaffolded groups. To correct for multiple comparisons, we used false discover rate (FDR) correction in the core *R* package *stats*. For the significant effects from the ANOVAs, we conducted one-tailed *t*-tests on change scores to evaluate directional hypotheses from the preregistration. We used one-tailed *t*-tests as we hypothesized larger decreases in the mindfulness groups than the audiobook groups (e.g. larger decreases in stress). Additionally, we used ANCOVA to evaluate group effects on changes in outcomes over time (i.e., post-test scores predicted by group covarying pre-test scores) (Supplementary Table S3). We used multiple imputation to handle missing data which is robust to data that are missing at random (Graham, 2009). Multiple imputation was implemented using the *mice* (van Buuren & Groothuis-Oudshoorn, 2011), *jomo* (Quartagno & Carpenter, 2020), and *mitools* (Lumley, 2019) packages in *R*. We conducted sensitivity analyses, assessing the pre-post effects using ANCOVA while also covarying demographic variables (Supplementary Table S4). We also explored whether demographic variables moderated the effect of group assignment for our main measures of anxiety and child perceived stress (Supplementary Table S5, S6). Lastly, to examine dosage effects in the mindfulness group, we correlated the changes in measures with

number of days practiced, and then conducted a median split by days practiced (a median was used because of deviations from normality in the distribution of days practiced; Supplementary Fig. S1). Dosage effects in the audiobook groups were assessed using minutes listened by each participant.

Note that there were deviations from our preregistered analysis plan. Firstly, we intended to test hypotheses about a foraging task, but were unable to do so due to programming issues. Secondly, we did not investigate mediation by hours practiced because that data was unavailable, and instead conducted correlations between the number of days practiced and pre-post changes in the measures. We assessed these dosage effects for all measures. The following analyses were not preregistered: one-tailed *t*-tests on change scores to examine the significant group X time effects from the ANOVAs, ANCOVAs controlling for baseline scores, sensitivity tests (covarying demographics in ANCOVAs for all outcomes), and moderation tests (assessing the interaction of demographics with group assignment for main outcomes).

## Results

As assessed by *Inner Explorer*, the Mindfulness group practiced on average 25.30 ( $SD = 12.19$ ) of the targeted 40 days (Supplementary Fig. S1). The most frequently selected option by parents for number of practices was 40 (Supplementary Fig. S2). *Inner Explorer* days practiced and parent-report number of practices were positively related (Supplementary Fig. S3). The Audiobook-only group listened to audiobooks on average for 839.16 minutes ( $SD = 1231.91$  min), and the Audiobook-scaffolded group listened to audiobooks on average for 1022.26 minutes ( $SD = 1000.89$  min).

### Attrition

Overall 84.6% (236/279) of children completed the intervention. Specifically, 85.6% of Mindfulness participants (77/90), 85.6% of Audiobook-only participants (77/90), and 82.8% of Audiobook-scaffolded participants (82/99) who completed pre-test also completed at least some post-test. There was no significant difference in completion of pre-test between the mindfulness and audiobook groups ( $p = 0.92$ ,  $OR = 0.96$ , 95% CI [0.44 2.00]). There was no significant difference in completion of the intervention between groups ( $p = 0.76$ ,  $OR = 0.89$ , 95% CI [0.43 1.78]). The only demographic variable associated with completion rates was maternal education; higher maternal education correlated with higher completion rates across all groups ( $p = 0.013$ ,  $OR = 2.61$ , 95% CI [1.12 6.58]). Further, completion was not associated with outcome variables at baseline (all  $p$ -values were greater than 0.15).

### Power Analyses

We conducted post-hoc power analyses using the *pwr.t2n.test* and *pwr.anova.test* functions in the *pwr* package in *R* (Champlsey, 2020). For comparisons of change scores between the mindfulness and audiobook conditions, we were powered at 80% to detect Cohen's  $d$ s of at least 0.42 at a significance level of  $p = 0.05$ . We also conducted a power analysis of the ANOVA between the three conditions, which revealed that we were powered at 80% to detect  $f$ -values of 0.19 at a significance level of  $p = 0.05$ .

----- Table 2 -----

### Outcome Analyses

Correlations between outcome measures at baseline can be found in Supplementary Table S1. The three groups did not differ significantly on demographic or outcome measures at baseline ( $ps \geq 0.05$ ; Table 1, 2), with one exception. Specifically, the Mindfulness group had a

higher proportion of child mental health diagnoses than the Audiobook-only group (Tukey test,  $p = 0.046$ ). We controlled for possible effects of this difference in a sensitivity analysis using ANCOVA (Supplementary Table S4).

-----Table 3-----

Effect sizes were generally small (Cohen's  $d$ s < 0.45, Table 3). When controlling for multiple comparisons, there was one significant group X time interaction using pairwise ANOVAs (Fig. 2, Table 3). Parent-reported child negative affect decreased significantly more in the Mindfulness group than the Audiobook-only group ( $F(1,146) = 8.60$ , FDR-corrected  $p = 0.048$ ). Child-perceived stress also decreased more in the Mindfulness group than the Audiobook-only group, although this trend was not significant ( $F(1,146) = 3.20$ , uncorrected  $p = 0.075$ ). No significant group X time interactions were observed for the comparison of Mindfulness vs Audiobook-scaffolded groups.

We examined these effects from the ANOVAs, calculating one-tailed  $t$ -tests between change scores for child-perceived stress and parent-reported negative affect because those measures yielded significant or trending towards significant differences. We found that reductions in child-perceived stress in the Mindfulness group were significantly greater than in the Audiobook-only group ( $t(148,74) = 1.79$ ,  $p = 0.038$ ). We also found that reductions in parent-reported negative affect in the Mindfulness group were significantly greater than in the Audiobook-only group ( $t(130,05) = 2.89$ ,  $p = 0.0022$ ).

Time effects across all interventions were observed for reduced child-reported negative affect, and also parent reports of improved global executive composite (GEC), improved behavioral regulation index (BRI), reduced child depression symptoms, reduced parent stress, and reduced child negative affect (Supplementary Table S2).

----- Fig. 2 -----

When controlling for multiple comparisons, there were no differences between groups on pre-post changes in main outcomes or exploratory outcomes using ANCOVA models (Table S3,  $p$ -values  $> 0.10$ ). However, there was a trend difference between Mindfulness and Audiobook-only groups (FDR-corrected  $p = 0.072$ ) in parent-reported child negative affect when covarying demographic variables, that is, children in the mindfulness group had greater decreases in parent-reported child negative affect (Supplementary Table S4). We further tested for possible baseline moderation by demographic variables and results were not significant (Supplementary Tables S5, S6).

We explored dosage effects using days practiced recorded by *Inner Explorer* in the Mindfulness group. We conducted a median split on days practiced where a Low-Practice group was defined as at or below the median (29 days), and a High-Practice group was defined as above the median. The High-Practice group had significantly larger decreases in parent-reported child negative affect than the Low-Practice group ( $t(63) = 2.07, p = 0.021$ ), and significantly larger decreases in parental stress ( $t(63) = 4.16, p < 0.001$ ). The High-Practice group had significantly larger decreases in parent-reported negative affect than both the Audiobook-scaffolded ( $t(115) = 2.18, p = 0.016$ ), and Audiobook-only ( $t(104) = 2.98, p = 0.0018$ ) groups (Fig. 3). The High-Practice group had significantly larger decreases in parental stress than both the Audiobook-scaffolded ( $t(115) = 2.53, p = 0.0065$ ), and Audiobook-only ( $t(104) = 3.16, p = 0.0010$ ) groups. The High-Practice group had significantly larger decreases in child-perceived stress than the Audiobook-only group ( $t(112) = 1.66, p = 0.049$ ), and trended toward larger decreases in child-perceived stress than the Audiobook-scaffolded group ( $t(117) = 1.32, p = 0.094$ ). Dosage effects were similar for parent reports of number of practices (Supplementary

Fig. S4). No dosage effects were found for all other outcomes. No dosage effects were found in the reading minutes for the audiobook groups when correlating minutes practiced with outcomes (parent-reported child negative affect  $p = 0.12$ , parental stress  $p = 0.18$ , child perceived stress  $p = 0.92$ ).

----- Fig. 3 -----

### Discussion

In this study, we administered an at-home app-based mindfulness intervention to children ages 8 to 10 years in the US. We hypothesized that there would be decreases in child-reported anxiety symptoms and perceived stress after the mindfulness intervention relative to the control audiobook interventions. There was evidence of larger decreases in children's negative affect as reported by parents and of children's self-perceived stress in the Mindfulness intervention compared to the Audiobook-only intervention, but no differences between the Mindfulness intervention and the Audiobook-scaffolded intervention. The reduction in stress is consistent with mindfulness studies reporting reduced stress with adults (Khoury et al., 2015) and children (Bauer et al., 2019). There were no significant group differences for trait mindfulness, parent-report anxiety and depression, executive function, prosociality, and parental stress. Effect sizes in all interventions were small (Cohen's  $d$ s < 0.45). There was evidence that participants who engaged with the mindfulness intervention more often showed larger reductions in negative affect.

This study was novel in applying a remote, app-based intervention with pre-adolescent children. The remote nature of the mindfulness intervention may have diminished its effectiveness. While app-based mindfulness interventions in adults often show decreases in anxiety and depression (Spijkerman et al., 2016), increases in well-being (Gál et al., 2021) and

increases in mindfulness (Linardon et al., 2019), changes are often smaller in size than in-person interventions (Goldberg et al., 2018). Indeed, this has been observed across app-based health interventions generally (Goldberg et al., 2022a). There are barriers to success for app-based interventions, including low adherence (Linardon & Fuller-Tyszkiewicz, 2020). However, in the current study we sought to encourage adherence by including weekly practice reminders as well as personal communication in the case of multiple missed practices. Our monitoring data show that most families were practicing (on average 25 days of a total of 40), and dropout was relatively low compared to other app-based studies (e.g., Goldberg et al., 2020; Linardon & Fuller-Tyszkiewicz, 2020).

Another potential explanation for the small effects is the rigor of the control conditions. It is well-established that active control conditions produce smaller effect sizes (Gál et al., 2021; Goldberg et al., 2022b). In our study, participants were recruited for a reading study to evaluate the impact of audiobooks on reading scores, and were randomized into either Audiobook-only, Audiobook-scaffolded (with weekly meetings with a facilitator), or Mindfulness conditions. For our mindfulness analyses, we thus had two audiobook control conditions. It is possible that these audiobook conditions were not the best comparison conditions for mindfulness, especially because the Audiobook-scaffolded condition involved weekly contact with a facilitator which was not present in the mindfulness condition. The interventions took place during the height of the COVID-19 pandemic in the US, when social and academic restrictions were in place. Given the circumstances, the human contact provided by the Audiobook-scaffolded intervention may have been particularly beneficial.

Indeed, there were multiple pre-post improvements across all three interventions, including measures of children's negative feelings, and parent ratings of children's negative

affect, depression symptoms, executive function, and behavior regulation. One possibility is that the human contact involved in remote but interactive pre-post testing, and the activity involved in all three interventions were beneficial in the context of COVID isolation. Social support and interaction was an important determiner of mental health during the COVID-19 pandemic for adults (Li 2021; Saltzman et al., 2020), and likely in children as well (Wong et al., 2020).

Whether all three interventions would show similar benefits in more typical circumstances is unknown. Additionally, it is possible that parents had biased expectancies for benefits for all the interventions.

Our study is not alone in finding limited effectiveness of mindfulness interventions for children. A school-based RCT with over 8,000 adolescents found that mindfulness interventions showed no advantages over teaching as usual (Montero-Marin et al., 2022). In addition, they found that the mindfulness interventions were actually contraindicated for some adolescents with baseline mental health difficulties. Other, smaller studies with children and adolescents have also found null results (Malboeuf-Hurtubise et al., 2021; Odgers et al., 2020). A meta-analysis of mindfulness interventions in youth found no increases in well-being relative to active or passive (i.e., waitlist or treatment-as-usual) controls, and only small decreases in anxiety and stress with regard to active controls (Dunning et al., 2022).

In contrast, numerous studies show evidence for the benefits of mindfulness in children. In child and adolescent populations, mindfulness-based interventions (MBIs) have been related to symptom reduction, increased wellbeing (Caldwell et al., 2019; Carsley et al., 2018; Porter et al., 2022), improved attention (Bauer et al., 2020), and behavioral regulation (Kaunhoven & Dorjee, 2017; Schonert-Reichl et al., 2015). Researchers have found functional changes in brain circuitry in sixth graders after a mindfulness intervention (Bauer et al., 2019). Trait mindfulness,

defined as the disposition or tendency to behave mindfully in day-to-day life, is connected to positive mental health outcomes in children (de Bruin et al., 2014; Greco et al., 2011) and positive academic outcomes and school behaviors (Caballero et al., 2019). In theory, MBIs promote trait mindfulness and thus positive mental health outcomes.

The benefit of the mindfulness app on children's emotional well-being (child perceived stress and parent-reported child negative affect) was related to frequency of the usage of the app. Children who practiced 30 days or more had significantly larger gains in emotional well-being than children in both the Audiobook-only and Audiobook-scaffolded conditions. We also found a similar dose-dependent benefit for parental stress. These results suggest there needs to be sufficient dosage for app-based interventions to show effects in children, paralleling similar findings in adults (Bostock et al., 2019). Further, the large school-based study of mindfulness in the UK that found no benefits contained only 10 instructional sessions spread across a semester and that most students reported little or no independent practice beyond those sessions (Montero-Marín et al., 2022). In contrast, for example, a school-based study that yielded multiple benefits with 6<sup>th</sup> graders involved about 24 hours of instruction and practice across 8 weeks (Bauer et al., 2019, 2020). It is, however possible that participants in the present study who practiced more than 30 days were more motivated to report mindfulness-related benefits. Future studies could randomize participants to receive more or less treatment so that there is direct evidence of the relation between dosage and possible benefits and so that the minimally effective dosage might be identified.

These findings suggest that an important issue in at-home app-based mindfulness interventions is adherence to the program. For young children, such adherence most likely involves adult caregivers as well as the children who depend on them. This challenge contrasts

with school-based programs in which, when given with high fidelity, the instructor can assure engagement.

### **Limitations and Future Research**

There are some limitations to our study that may account for the limited effectiveness of the mindfulness intervention. One limitation was that the families were initially recruited to take part in an audiobook intervention. Thus, they may have been less engaged and motivated to take part in the mindfulness intervention. There is evidence that creating more favorable subjective norms around mindfulness and increasing intentions by emphasizing the benefits of mindfulness (e.g. behavioral control) increases the number of minutes practiced by participants (Crandall et al., 2019). Initial motivations to engage in mindfulness practice also appear to be related to persistence (Jiwani et al., 2022), and motivations for mindfulness may have been initially low in this sample. As participants were informed about their intervention assignment before pre-test, any difference in motivation could have effects not just on adherence, but also on test scores. However, we found no baseline differences in scores between the groups, no differences in completion of pre-testing, and no differences in completion of the interventions.

An additional limitation is that we may have been underpowered to detect small to moderate selective effects between interventions. Power analyses for interventions have been conducted using between-group comparisons of change scores (Goldberg et al., 2020), or  $f$ -values (Kaplan et al., 2022). We were powered at 80% to detect effect sizes of  $d = 0.42$  or larger for the comparisons of Mindfulness and Audiobook groups. We were powered at 80% to detect  $f$ -values of 0.19 using the ANOVA analyses. Future studies will ideally be more highly powered, but there may be questions about value of benefits that are so small that large numbers of participants are needed to detect them. There is the possibility, however, that the individual

differences among participants are predictive of the magnitude of benefits (Webb et al., 2022). Also, to better generalize to the public, studies should be conducted with diverse participants. While we focused substantial effort and resources toward recruiting a socioeconomically diverse sample (Ozernov-Palchik et al., 2022) the majority of participants in our study were from middle to high-income families.

The final limitation concerns the validity of our self-report measures. We used a clinical questionnaire (the RCADS (Chorpita et al., 2000)) to measure anxiety symptoms. Floor effects may have obscured sub-clinical changes in anxiety. In addition, the child perceived stress measure had low scale reliability.

In summary, we conducted a rigorously controlled, app-based mindfulness intervention in children, and found some limited benefits in reduced stress and enhanced emotional well-being. Smartphone apps offer benefits in terms of scalability, reach, and cost-effectiveness (Linardon & Fuller-Tyszkiewicz, 2020). For this reason, we suggest that future work should continue to study remote interventions in children, examining the types of mindfulness practices that are useful, how much practice is necessary to achieve beneficial outcomes, how to encourage children to accomplish that amount of practice, and which children may benefit most from a mindfulness app.

## **Declarations**

### **Compliance with Ethical Standards**

This project received Institutional Review Board approval at the Massachusetts Institute of Technology (MIT).

**Consent**

We obtained informed consent from the parents of all participants as well as assent from the children before they participated in the study.

**Conflicts Of Interest**

The authors have no relevant financial or non-financial interests to disclose.

**Funding Statement**

This research was funded by the Chan Zuckerberg Initiative as part of the Reach Every Reader Project, National Institutes of Health (F32-HD100064 to OO), and the National Science Foundation Graduate Research Fellowship (Grant No. 1745302 to HO). SBG was supported by the National Center for Complementary & Integrative Health of the National Institutes of Health under Award Number K23AT010879.

**Data, Materials and Code Availability**

Data and code will be made publicly available on publication at <https://osf.io/a9h6t/>

**Author Contributions**

Isaac Treves: Conceptualization, analysis, writing-original draft preparation, supervision. Halie Olson: Conceptualization, project administration, funding acquisition, writing-review and editing. Ola Ozernov-Palchik: Conceptualization, project administration, funding acquisition, supervision. Cindy Li: data curation, writing-original draft preparation, investigation, methodology. Kimberly Wang: project administration, writing-review and editing, investigation.

Xochitl Arechiga: project administration, investigation. Simon Goldberg: analysis, writing-review and editing. John Gabrieli: supervision, resources, writing-review and editing.

### **Use of Artificial Intelligence**

Artificial Intelligence was not used.

### **Acknowledgements**

We thank our testers: Amanda Miller, David Bates, Ross Weissman, Joohee Baik, June Okada, William Oliver, and Harriet Richards. We also thank Yesi Camacho Torres, Hope Kentala, Natalie Gardino, and Jeff Dieffenbach for helping to run the study. We thank the Los Angeles Unified School District (LAUSD) Achievement Network for help with recruiting families. We thank Learning Ally for help with recruitment and for providing the audiobook interventions. Lastly, we thank Inner Explorer for providing the intervention.

### **References**

- American Psychiatric Association. (1994). *Diagnostic and statistical manual of mental disorders* (4th ed.).
- Anderson, M., & Jiang, J. (2018). *Teens, Social Media and Technology 2018*. Pew Research Center. Retrieved July 14, 2023, from <https://www.pewresearch.org/internet/2018/05/31/teens-social-media-technology-2018/>
- Bakosh, L. S. (2013). *Investigating the effects of a daily audio-guided mindfulness intervention for elementary school students and teachers* (Publication No. 3618722) [Doctoral

- dissertation, Sofia University]. ProQuest Dissertations Publishing.
- Bakosh, L. S., Snow, R. M., Tobias, J. M., Houlihan, J. L., & Barbosa-Leiker, C. (2016). Maximizing mindful learning: Mindful awareness intervention improves elementary school students' quarterly grades. *Mindfulness*, 7(1), 59–67. <https://doi.org/10.1007/s12671-015-0387-6>
- Bauer, C. C. C., Caballero, C., Scherer, E., West, M. R., Mrazek, M. D., Phillips, D. T., Whitfield-Gabrieli, S., & Gabrieli, J. D. E. (2019). Mindfulness training reduces stress and amygdala reactivity to fearful faces in middle-school children. *Behavioral Neuroscience*, 133(6), 569–585. <https://doi.org/10.1037/bne0000337>
- Bauer, C. C. C., Rozenkrantz, L., Caballero, C., Nieto-Castanon, A., Scherer, E., West, M. R., Mrazek, M., Phillips, D. T., Gabrieli, J. D. E., & Whitfield-Gabrieli, S. (2020). Mindfulness training preserves sustained attention and resting state anticorrelation between default-mode network and dorsolateral prefrontal cortex: A randomized controlled trial. *Human Brain Mapping*, 41(18), 5356–5369. <https://doi.org/10.1002/hbm.25197>
- Beauchaine, T. P., Gatzke-Kopp, L., & Mead, H. K. (2007). Polyvagal theory and developmental psychopathology: Emotion dysregulation and conduct problems from preschool to adolescence. *Biological Psychology*, 74(2), 174-184. <https://doi.org/10.1016/j.biopsycho.2005.08.008>
- Bostock, S., Crosswell, A. D., Prather, A. A., & Steptoe, A. (2019). Mindfulness on-the-go: Effects of a mindfulness meditation app on work stress and well-being. *Journal of Occupational Health Psychology*, 24(1), 127–138. <https://doi.org/10.1037/ocp0000118>
- Caballero, C., Scherer, E., West, M. R., Mrazek, M. D., Gabrieli, C. F. O., & Gabrieli, J. D. E.

- (2019). Greater mindfulness is associated with better academic achievement in middle school. *Mind, Brain, and Education*, 13(3), 157-166. <https://doi.org/10.1111/mbe.12200>
- Caldwell, D. M., Davies, S. R., Hetrick, S. E., Palmer, J. C., Caro, P., López-López, J. A., Gunnell, D., Kidger, J., Thomas, J., French, C., Stockings, E., Campbell, R., & Welton, N. J. (2019). School-based interventions to prevent anxiety and depression in children and young people: A systematic review and network meta-analysis. *Lancet Psychiatry*, 6(12), 1011–1020. [https://doi.org/10.1016/S2215-0366\(19\)30403-1](https://doi.org/10.1016/S2215-0366(19)30403-1)
- Carsley, D., Khoury, B., & Heath, N. L. (2018). Effectiveness of mindfulness interventions for mental health in schools: A comprehensive meta-analysis. *Mindfulness*, 9(3), 693–707. <https://doi.org/10.1007/s12671-017-0839-2>
- Centers for Disease Control and Prevention. (2022). *Data and statistics on children's mental health*. <https://www.cdc.gov/childrensmentalhealth/data.html>
- Champley, S. (2020). *pwr: Basic functions for power analysis*. <https://cran.r-project.org/web/packages/pwr/pwr.pdf>
- Chorpita, B. F., Yim, L., Moffitt, C. E., Umemoto L. A., & Francis, S. E. (2000). Assessment of symptoms of DSM-IV anxiety and depression in children: A revised child anxiety and depression scale. *Behaviour Research and Therapy*, 38(8), 835-855. [https://doi.org/10.1016/S0005-7967\(99\)00130-8](https://doi.org/10.1016/S0005-7967(99)00130-8)
- Cohen, S., Kamarck, T., & Mermelstein, R. (1983). A global measure of perceived stress. *Journal of Health and Social Behavior*, 24(4), 385–396. <https://doi.org/10.2307/2136404>
- Cohen, S. (1988). Perceived stress in a probability sample of the United States. In S. Spacapan & S. Oskamp (Eds.), *The social psychology of health* (pp. 31–67). Sage Publications, Inc.

- Crandall, A., Cheung, A., Young, A., & Hooper, A. P. (2019). Theory-based predictors of mindfulness meditation mobile app usage: A survey and cohort study. *JMIR MHealth and UHealth*, 7(3), 1–11. <https://doi.org/10.2196/10794>
- Creswell, J. D. (2017). Mindfulness interventions. *Annual Review of Psychology*, 68(1), 491-516. <https://doi.org/10.1146/annurev-psych-042716-051139>
- Davidson, R. J., & Kaszniak, A. W. (2015). Conceptual and methodological issues in research on mindfulness and meditation. *American Psychologist*, 70(7), 581–592. <https://doi.org/10.1037/a0039512>
- de Bruin, E. I., Zijlstra, B. J. H., & Bögels, S. M. (2014). The meaning of mindfulness in children and adolescents: Further validation of the child and adolescent mindfulness measure (CAMM) in two independent samples from the Netherlands. *Mindfulness*, 5, 422-430. <https://doi.org/10.1007/s12671-013-0196-8>
- Diamond, A., & Lee, K. (2011). Interventions shown to aid executive function development in children 4-12 years old. *Science*, 333(6045), 959-964. <https://doi.org/10.1126/science.1204529>
- Dunning, D., Tudor, K., Radley, L., Dalrymple, N., Funk, J., Vainre, M., Ford, T., Montero-Marín, J., Kuyken, W., & Dalgleish, T. (2022). Do mindfulness- based programmes improve the cognitive skills, behaviour and mental health of children and adolescents? An updated meta-analysis of randomised controlled trials. *Evidence Based Mental Health*, 25(3), 135–142. <https://doi.org/10.1136/ebmental-2022-300464>
- Dunlap, R. D. (2023). *An Analysis of the Impact of the Inner Explorer Mindfulness Program on the Academic Growth of Elementary Students* (Doctoral dissertation, The University of North Carolina at Charlotte).

- Ebesutani, C., Reise, S. P., Chorpita, B. F., Ale, C., Regan, J., Young, J., Higa-McMillan, C., & Weisz, J. R. (2012). The revised child anxiety and depression scale-short version: Scale reduction via exploratory bifactor modeling of the broad anxiety factor. *Psychological Assessment, 24*(4), 833-845. <https://doi.org/10.1037/a0027283>
- Ebesutani, C., Korathu-Larson, P., Nakamura, B. J., Higa-McMillan, C., & Chorpita, B. (2017). The revised child anxiety and depression scale 25-parent version: Scale development and validation in a school-based and clinical sample. *Assessment, 24*(6), 712–728. <https://doi.org/10.1177/1073191115627012>
- Economides, M., Martman, J., Bell, M. J., & Sanderson, B. (2018). Improvements in stress, affect, and irritability following brief use of a mindfulness-based smartphone app: A randomized controlled trial. *Mindfulness, 9*(5), 1584–1593. <https://doi.org/10.1007/s12671-018-0905-4>
- Emerson, C. S., Mollet, G. A., & Harrison, D. W. (2005). Anxious-depression in boys: An evaluation of executive functioning. *Archives of Clinical Neuropsychology, 20*(4), 539-546. <https://doi.org/10.1016/j.acn.2004.10.003>
- Flett, J. A. M., Hayne, H., Riordan, B. C., Thompson, L. M., & Conner, T. S. (2019). Mobile mindfulness meditation: A randomised controlled trial of the effect of two popular apps on mental health. *Mindfulness, 10*(5), 863–876. <https://doi.org/10.1007/s12671-018-1050-9>
- Flook, L., Goldberg, S. B., Pinger, L., & Davidson, R. J. (2015). Promoting prosocial behavior and self-regulatory skills in preschool children through a mindfulness-based kindness curriculum. *Developmental Psychology, 51*(1), 44-51. <https://doi.org/10.1037/a0038256>
- Friedman, N. P., Haberstick, B. C., Willcutt, E. G., Miyake, A., Young, S. E., Corley, R. P., &

- Hewitt, J. K. (2007). Greater attention problems during childhood predict poorer executive functioning in late adolescence. *Psychological Science, 18*(10), 893-900.  
<https://doi.org/10.1111/j.1467-9280.2007.01997.x>
- Gál, É., Ștefan, S., & Cristea, I. A. (2021). The efficacy of mindfulness meditation apps in enhancing users' well-being and mental health related outcomes: A meta-analysis of randomized controlled trials. *Journal of Affective Disorders, 279*, 131–142.  
<https://doi.org/10.1016/j.jad.2020.09.134>
- Gioia, G. A., Isquith, P. K., Guy, S. C., & Kenworthy, L. (2000). Test review: Behavior rating inventory of executive function. *Child Neuropsychology, 6*(3), 235-238.  
<https://doi.org/10.1076/chin.6.3.235.3152>
- Goldberg, S. B., Imhoff-Smith, T., Bolt, D. M., Wilson-Mendenhall, C. D., Dahl, C. J., Davidson, R. J., & Rosenkranz, M. A. (2020). Testing the efficacy of a multicomponent, self-guided, smartphone-based meditation app: Three-armed randomized controlled trial. *JMIR Mental Health, 7*(11). <https://doi.org/10.2196/23825>
- Goldberg, S. B., Lam, S. U., Simonsson, O., Torous, J., & Sun, S. (2022). Mobile phone-based interventions for mental health: A systematic meta-review of 14 meta-analyses of randomized controlled trials. *PLOS Digital Health, 1*(1).  
<https://doi.org/10.1371/journal.pdig.0000002>
- Goldberg, S. B., Riordan, K. M., Sun, S., & Davidson, R. J. (2022). The empirical status of mindfulness-based interventions: A systematic review of 44 meta-analyses of randomized controlled trials. *Perspectives on Psychological Science, 17*(1), 108–130.  
<https://doi.org/10.1177/1745691620968771>
- Goldberg, S. B., Tucker, R. P., Greene, P. A., Davidson, R. J., Wampold, B. E., Kearney, D. J.,

- & Simpson, T. L. (2018). Mindfulness-based interventions for psychiatric disorders: A systematic review and meta-analysis. *Clinical Psychology Review, 59*, 52-60.  
<https://doi.org/10.1016/j.cpr.2017.10.011>
- Goodman, M. S. (2019). *Targeting self-regulation and disease resilience in elementary school students through a mindfulness-based social-emotional learning curriculum* (Publication No. 10826744) [Doctoral dissertation, Alliant International University]. ProQuest Dissertations Publishing.
- Graham, J. W. (2009). Missing data analysis: Making it work in the real world. *Annual Review of Psychology, 60*, 549–576. <https://doi.org/10.1146/annurev.psych.58.110405.085530>
- Graziano, P. A., Reavis, R. D., Keane, S. P., & Calkins, S. D. (2007). The role of emotion regulation in children's early academic success. *Journal of School Psychology, 45*(1), 3-19. <https://doi.org/10.1016/j.jsp.2006.09.002>
- Greco, L. A., Baer, R. A., & Smith, G. T. (2011). Assessing mindfulness in children and adolescents: Development and validation of the child and adolescent mindfulness measure (CAMM). *Psychological Assessment, 23*(3), 606–614.  
<https://doi.org/10.1037/a0022819>
- Greenberg, M. T., & Harris, A. R. (2012). Nurturing mindfulness in children and youth: Current state of research. *Child Development Perspectives, 6*(2), 161–166.  
<https://doi.org/10.1111/j.1750-8606.2011.00215.x>
- Greenland, S. K. (2010). *The mindful child: How to help your kid manage stress and become happier, kinder, and more compassionate*. Simon and Schuster.
- Grist, R., Croker, A., Denne, M., & Stallard, P. (2019). Technology delivered interventions for depression and anxiety in children and adolescents: A systematic review and meta-

analysis. *Clinical Child and Family Psychology Review*, 22(2), 147–171.

<https://doi.org/10.1007/s10567-018-0271-8>

Guhn, M., Gadermann, A. M., Almas, A., Schonert-Reichl, K. A., & Hertzman, C. (2016).

Associations of teacher-rated social, emotional, and cognitive development in kindergarten to self-reported wellbeing, peer relations, and academic test scores in middle childhood. *Early Childhood Research Quarterly*, 35, 76-84.

<https://doi.org/10.1016/j.ecresq.2015.12.027>

Harris, P. A., Taylor, R., Thielke, R., Payne, J., Gonzalez, N., & Conde, J. G. (2009). Research

electronic data capture (REDCap)—A metadata-driven methodology and workflow process for providing translational research informatics support. *Journal of Biomedical Informatics*, 42(2), 377–381.

<https://doi.org/10.1016/J.JBI.2008.08.010>

Hilt, L. M., & Swords, C. M. (2021). Acceptability and preliminary effects of a mindfulness

mobile application for ruminative adolescents. *Behavior Therapy*, 52(6), 1339–1350.

<https://doi.org/10.1016/j.beth.2021.03.004>

Hilt, L. M., Swords, C. M., & Webb, C. A. (2023). Randomized controlled trial of a mindfulness

mobile application for ruminative adolescents. *Journal of Clinical Child & Adolescent Psychology*, 00(00), 1–14. <https://doi.org/10.1080/15374416.2022.2158840>

Hirshberg, M. J., Frye, C., Dahl, C. J., Riordan, K. M., Vack, N. J., Sachs, J., Goldman, R.,

Davidson, R. J., & Goldberg, S. B. (2022). A randomized controlled trial of a smartphone-based well-being training in public school system employees during the COVID-19 pandemic. *Journal of Educational Psychology*, 114(8), 1895–1911.

<https://doi.org/10.1037/edu0000739>

Inner Explorer. (2023). *Inner Explorer* (Version 7.5.3) [Mobile app]. Google Play Store.

[https://play.google.com/store/apps/details?id=com.innerexplorer.present&hl=en\\_US  
gl=US&pli=1](https://play.google.com/store/apps/details?id=com.innerexplorer.present&hl=en_US&gl=US&pli=1)

Jiwani, Z., Lam, S. U., Davidson, R. J., & Goldberg, S. B. (2022). Motivation for meditation and its association with meditation practice in a national sample of internet users.

*Mindfulness*, 13(10), 2641–2651. <https://doi.org/10.1007/s12671-022-01985-6>

Kabat-Zinn, J. (2003). Mindfulness-based interventions in context: Past, present, and future. *Clinical Psychology: Science and Practice*, 10(2), 144-156.

<https://doi.org/10.1093/clipsy.bpg016>

Kabat-Zinn, J. (2005). *Full catastrophe living: Using the wisdom of your body and mind to face stress, pain, and illness* (15th anniversary ed.). Delta Trade Paperback/Bantam Dell.

Kaplan, D. M., Mehl, M. R., Pace, T. W. W., Negi, L. T., Silva, B. O. de, Lavelle, B. D., Sivilli, T., Williams, A., Comstock, T., Price, B., Medrano, V., Robbins, M. L., Cole, S. P.,

Craighead, W. E., & Raison, C. L. (2022). Implications of a “null” randomized controlled trial of mindfulness and compassion interventions in healthy adults. *Mindfulness*, 13(5), 1197–1213. <https://doi.org/10.1007/s12671-022-01861-3>

Kaufman, A. S. (2004). *Kaufman brief intelligence test—second edition (KBIT-2)*. Circle Pines, MN: American Guidance Service.

Kaunhoven, R. J., & Dorjee, D. (2017). How does mindfulness modulate self-regulation in pre-adolescent children? An integrative neurocognitive review. *Neuroscience and*

*Biobehavioral Reviews*, 74, 163–184. <https://doi.org/10.1016/j.neubiorev.2017.01.007>

Khoury, B., Sharma, M., Rush, S. E., & Fournier, C. (2015). Mindfulness-based stress reduction for healthy individuals: A meta-analysis. *Journal of Psychosomatic Research*, 78(6), 519-528. <https://doi.org/10.1016/j.jpsychores.2015.03.009>

Learning Ally. (2023). *Learning Ally* (Version 4.7.3) [Mobile app]. Google Play Store.

[https://play.google.com/store/apps/details?id=org.learningally.LinkMobile&hl=en\\_US&gl=US](https://play.google.com/store/apps/details?id=org.learningally.LinkMobile&hl=en_US&gl=US)

Li, X., Fu, P., Fan, C., Zhu, M., & Li, M. (2021). COVID-19 stress and mental health of students in locked-down colleges. *International Journal of Environmental Research and Public Health*, *18*(2), 771. <https://doi.org/10.3390/ijerph18020771>

Linardon, J., Cuijpers, P., Carlbring, P., Messer, M., & Fuller-Tyszkiewicz, M. (2019). The efficacy of app-supported smartphone interventions for mental health problems: A meta-analysis of randomized controlled trials. *World Psychiatry*, *18*(3), 325–336.

<https://doi.org/10.1002/wps.20673>

Linardon, J., & Fuller-Tyszkiewicz, M. (2020). Attrition and adherence in smartphone-delivered interventions for mental health problems: A systematic and meta-analytic review. *Journal of Consulting and Clinical Psychology*, *88*(1), 1–13. <https://doi.org/10.1037/ccp0000459>

Liverpool, S., Mota, C. P., Sales, C. M. D., Čuš, A., Carletto, S., Hancheva, C., Sousa, S., Cerón, S. C., Moreno-Peral, P., Pietrabissa, G., Moltrecht, B., Ulberg, R., Ferreira, N., & Edbrooke-Childs, J. (2020). Engaging children and young people in digital mental health interventions: Systematic review of modes of delivery, facilitators, and barriers. *Journal of Medical Internet Research*, *22*(6). <https://doi.org/10.2196/16317>

Lumley, T. (2019). *mitools: Tools for multiple imputation of missing data*.

<https://cran.rproject.org/web/packages/mitools/mitools.pdf>

Lupien, S. J., McEwen, B. S., Gunnar, M. R., & Heim, C. (2009). Effects of stress throughout the lifespan on the brain, behaviour and cognition. *Nature Reviews Neuroscience*, *10*(6), 434–445. <https://doi.org/10.1038/nrn2639>

- Lutz, A., Slagter, H. A., Dunne, J. D., & Davidson, R. J. (2008). Attention regulation and monitoring in meditation. *Trends in Cognitive Sciences*, *12*(4), 163-169.  
<https://doi.org/10.1016/j.tics.2008.01.005>
- Mahlo, L., & Windsor, T. D. (2021). Feasibility, acceptability, and preliminary efficacy of an app-based mindfulness-meditation program among older adults. *The Gerontologist*, *61*(5), 775–786. <https://doi.org/10.1093/geront/gnaa093>
- Malboeuf-Hurtubise, C., Léger-Goodes, T., Mageau, G. A., Joussemet, M., Herba, C., Chadi, N., Lefrançois, D., Camden, C., Bussi eres,  . L., Taylor, G.,  thier, M. A., & Gagnon, M. (2021). Philosophy for children and mindfulness during COVID-19: Results from a randomized cluster trial and impact on mental health in elementary school students. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, *107*.  
<https://doi.org/10.1016/j.pnpbp.2021.110260>
- Mani, M., Kavanagh, D. J., Hides, L., & Stoyanov, S. R. (2015). Review and evaluation of mindfulness-based iPhone apps. *JMIR MHealth and UHealth*, *3*(3).  
<https://doi.org/10.2196/mhealth.4328>
- Miller, G. E., Chen, E., & Parker, K. J. (2011). Psychological stress in childhood and susceptibility to the chronic diseases of aging: Moving toward a model of behavioral and biological mechanisms. *Psychological Bulletin*, *137*(6), 959-997.  
<https://doi.org/10.1037/a0024768>
- Montero-Marin, J., Allwood, M., Ball, S., Crane, C., De Wilde, K., Hinze, V., Jones, B., Lord, L., Nuthall, E., Raja, A., Taylor, L., Tudor, K., Blakemore, S.-J., Byford, S., Dalglish, T., Ford, T., Greenberg, M. T., Ukoumunne, O. C., Williams, J. M. G., & Kuyken, W. (2022). School-based mindfulness training in early adolescence: What works, for whom

- and how in the MYRIAD trial? *Evidence Based Mental Health*, 25(3), 117–124.  
<https://doi.org/10.1136/EBMENTAL-2022-300439>
- Mrnjaus, K., & Krneta, M. (2014). Mindfulness, concentration and student achievement – Challenges and solutions. *Procedia-Social and Behavioral Sciences*, 116(21), 1044-1049.  
<https://doi.org/10.1016/j.sbspro.2014.01.343>
- Nunes, A., Castro, S. L., & Limpo, T. (2020). A review of mindfulness-based apps for children. *Mindfulness*, 11(9), 2089–2101. <https://doi.org/10.1007/s12671-020-01410-w>
- Odgers, K., Dargue, N., Creswell, C., Jones, M. P., & Hudson, J. L. (2020). The limited effect of mindfulness-based interventions on anxiety in children and adolescents: A meta-analysis. *Clinical Child and Family Psychology Review*, 23(3), 407–426.  
<https://doi.org/10.1007/s10567-020-00319-z>
- Ozernov-Palchik, O., Olson, H. A., Arechiga, X. M., Kentala, H., Solorio-Fielder, J. L., Wang, K. L., Torres, Y. C., Gardino, N. D., Dieffenbach, J. R., & Gabrieli, J. D. E. (2022). Implementing remote developmental research: A case study of a randomized controlled trial language intervention during COVID-19. *Frontiers in Psychology*, 12, 734375.  
<https://doi.org/10.3389/fpsyg.2021.734375>
- Panorama Education. (2015). *User guide: Panorama social-emotional learning survey*.  
<https://www.panoramaed.com/social-emotional-learning-sel>
- Polit, D. F., & Gillespie, B. M. (2010). Intention-to-treat in randomized controlled trials: Recommendations for a total trial strategy. *Research in Nursing & Health*, 33(4), 355-368. <https://doi.org/10.1002/nur.20386>
- Porter, B., Oyanadel, C., Sáez-Delgado, F., Andaur, A., & Peñate, W. (2022). Systematic review of mindfulness-based interventions in child-adolescent population: A developmental

- perspective. *European Journal of Investigation in Health, Psychology and Education*, 12(8), 1220–1243. <https://doi.org/10.3390/ejihpe12080085>
- Posner, M. I. (Ed.). (2011). *Cognitive neuroscience of attention*. The Guilford Press.
- Puzia, M., Laird, B., Green, J., & Huberty, J. (2020). Parents' perceptions of their children's engagement in a consumer-based meditation mobile app: Cross-sectional survey study. *JMIR Pediatrics and Parenting*, 3(2), 1–8. <https://doi.org/10.2196/24536>
- Quartagno, M., Grund, S., & Carpenter, J. (2020). Jomo: A flexible package for two-level joint modelling multiple imputation. *R Journal*, 9(1).
- Repetti, R. L., Taylor, S. E., & Seeman, T. E. (2002). Risky families: Family social environments and the mental and physical health of offspring. *Psychological Bulletin*, 128(2), 330-366. <https://doi.org/10.1037/0033-2909.128.2.330>
- Rideout, V. (2015). *The common sense census: Media use by tweens and teens*. <https://doi.org/10.3886/ICPSR38018.v1>
- Rideout, V., & Fox, S. (2018). *Digital health practices, social media use, and mental well-being among teens and young adults in the US*. <https://digitalcommons.psjhealth.org/cgi/viewcontent.cgi?article=2092&context=publications>
- Rotenberg, K. J., Michalik, N., Eisenberg, N., & Betts, L. R. (2008). The relations among young children's peer-reported trustworthiness, inhibitory control, and preschool adjustment. *Early Childhood Research Quarterly*, 23(2), 288-298. <https://doi.org/10.1016/j.ecresq.2007.04.003>
- Saltzman, L. Y., Hansel, T. C., & Bordnick, P. S. (2020). Loneliness, isolation, and social support factors in post-COVID-19 mental health. *Psychological Trauma: Theory*,

- Research, Practice, and Policy*, 12(S1), S55–S57. <https://doi.org/10.1037/tra0000703>
- Schonert-Reichl, K. A., Oberle, E., Lawlor, M. S., Abbott, D., Thomson, K., Oberlander, T. F., & Diamond, A. (2015). Enhancing cognitive and social–emotional development through a simple-to-administer mindfulness-based school program for elementary school children: A randomized controlled trial. *Developmental Psychology*, 51(1), 52–66. <https://doi.org/10.1037/a0038454>
- Schonfeld, D. J., Adams, R. E., Fredstrom, B. K., Weissberg, R. P., Gilman, R., Voyce, C., Tomlin, R., & Speese-Linehan, D. (2015). Cluster-randomized trial demonstrating impact on academic achievement of elementary social-emotional learning. *School Psychology Quarterly*, 30(3), 406–420. <https://doi.org/10.1037/spq0000099>
- Semple, R. J., Droutman, V., & Reid, B. A. (2017). Mindfulness goes to school: Things learned (so far) from research and real-world experiences. *Psychology in the Schools*, 54(1), 29–52. <https://doi.org/10.1002/pits.21981>
- Semple, R. J., Lee, J., Rosa, D., & Miller, L. F. (2010). A randomized trial of mindfulness-based cognitive therapy for children: Promoting mindful attention to enhance social-emotional resiliency in children. *Journal of Child and Family Studies*, 19, 218–229. <https://doi.org/10.1007/s10826-009-9301-y>
- Spijkerman, M. P. J., Pots, W. T. M., & Bohlmeijer, E. T. (2016). Effectiveness of online mindfulness-based interventions in improving mental health: A review and meta-analysis of randomised controlled trials. *Clinical Psychology Review*, 45, 102–114. <https://doi.org/10.1016/j.cpr.2016.03.009>
- Spinrad, T. L., Eisenberg, N., Cumberland, A., Fabes, R. A., Valiente, C., Shepard, S. A., Reiser, M., Losoya, S. H., & Guthrie, I. K. (2006). Relation of emotion-related regulation to

- children's social competence: A longitudinal study. *Emotion*, 6(3), 498-510.  
<https://doi.org/10.1037/1528-3542.6.3.498>
- Strickland, K. (2023). The effectiveness of utilizing the social-emotional learning program Inner Explorer on reducing disruptive behaviors in a Title 1 Setting [Doctoral dissertation, Saint Leo University].
- Thompson, R. A. (1994). Emotion regulation: A theme in search of definition. *Monographs of the Society for Research in Child Development*, 59(2-3), 25–52, 250-283.  
<https://doi.org/10.2307/1166137>
- Treves, I. N., Li, C. E., Wang, K. L., Ozernov-Palchik, O., Olson, H. A., & Gabrieli, J. D. E. (2023). Mindfulness supports emotional resilience in children during the COVID-19 pandemic. *PloS one*, 18(7), e0278501. <https://doi.org/10.1371/journal.pone.0278501>
- Tunney, C., Cooney, P., Coyle, D., & O'Reilly, G. (2017). Comparing young people's experience of technology-delivered v. face-to-face mindfulness and relaxation: Two-armed qualitative focus group study. *British Journal of Psychiatry*, 210(4), 284–289.  
<https://doi.org/10.1192/bjp.bp.115.172783>
- Turner, T., & Hingle, M. (2017). Evaluation of a mindfulness-based mobile app aimed at promoting awareness of weight-related behaviors in adolescents: A pilot study. *JMIR Research Protocols*, 6(4). <https://doi.org/10.2196/resprot.6695>
- Van Buuren, S., & Groothuis-Oudshoorn, K. (2011). mice: Multivariate imputation by chained equations in R. *Journal of Statistical Software*, 45(3), 1-67.  
<https://doi.org/10.18637/jss.v045.i03>
- Vickery, C. E., & Dorjee, D. (2016). Mindfulness training in primary schools decreases negative affect and increases meta-cognition in children. *Frontiers in Psychology*, 6, 1–13.

<https://doi.org/10.3389/fpsyg.2015.02025>

Webb, C. A., Hirshberg, M. J., Davidson, R. J., & Goldberg, S. B. (2022). Personalized prediction of response to smartphone-delivered meditation training: Randomized controlled trial. *Journal of Medical Internet Research*, 24(11), e41566.

<https://doi.org/10.2196/41566>

White, B. P. (2014). The perceived stress scale for children: A pilot study in a sample of 153 children. *International Journal of Pediatrics and Child Health*, 2, 45-52.

<https://doi.org/10.12974/2311-8687.2014.02.02.4>

Wong, C. A., Ming, D., Maslow, G., & Gifford, E. J. (2020). Mitigating the impacts of the COVID-19 pandemic response on at-risk children. *Pediatrics*, 146(1).

<https://doi.org/10.1542/peds.2020-0973>

**Table 1**

*Demographics of the Total Sample*

Variable	Overall ( <i>n</i> =279) % / mean ( <i>SD</i> )	Aud. + S. ( <i>n</i> =99) % / mean ( <i>SD</i> )	Aud. ( <i>n</i> =90) % / mean ( <i>SD</i> )	Mind. ( <i>n</i> =90) % / mean ( <i>SD</i> )	<i>p</i>
<b>Child Race/ethnicity</b>					
White	47.67	43.43	45.56	54.44	0.33
Black	2.15	0	5.56	1.11	
Latinx	10.75	13.13	11.11	7.78	
Asian	8.60	9.09	8.89	7.78	
Multiracial	22.58	23.23	22.22	22.22	
Not reported	8.24	11.11	6.67	6.67	
<b>Child Gender</b>					
Female	43.73	45.45	40.00	45.56	0.80
Male	56.27	54.55	60.00	54.44	
<b>Family Income</b>					
Less than \$20K	2.15	1.01	1.11	4.44	0.12
\$20-40K	5.38	9.09	5.56	1.11	
\$40-60K	7.17	7.07	8.89	5.56	
\$60-80K	10.75	10.10	15.56	6.67	
\$80-120K	20.43	22.22	20.00	18.89	

\$120-160K	13.62	12.12	12.22	16.67	
More than \$160K	31.54	26.26	30.00	38.89	
Not reported	8.96	12.12	6.67	7.78	
<b>Maternal Education</b>					0.88
7 <sup>th</sup> grade or less	0.36	0	1.11	0	
8 <sup>th</sup> /9 <sup>th</sup> grade	0.36	1.01	0	0	
10 <sup>th</sup> /11 <sup>th</sup> grade	1.43	1.01	2.22	1.11	
H.S.graduate/G.E.D.	10.04	13.13	10.00	6.67	
Associate Degree	8.96	8.08	12.22	6.67	
Bachelor Degree	24.37	22.22	22.22	28.89	
Master/Doctorate Degree	47.31	45.45	46.67	50.00	
Not reported	7.17	9.09	5.56	6.67	
<b>Child Age</b>	9.38 (0.54)	9.33 (0.58)	9.37 (0.48)	9.44 (0.55)	0.58
Not reported	0.72	1.01	0	1.11	
<b>Child Diagnosis* % (n)</b>	24.73 (69)	21.21 (21)	18.89 (17)	34.44 (31)	0.038
<b>Anxiety only</b>	4.66 (13)	2.02 (2)	3.33 (3)	8.89 (8)	
<b>ADHD only</b>	16.49 (46)	16.16 (16)	11.11 (10)	22.22 (20)	
<b>Anxiety &amp; ADHD</b>	3.58 (10)	3.03 (3)	4.44 (4)	3.33 (3)	

Note: Aud. + S. = Audiobook + Scaffolding Condition; Aud. = Audiobook Only Condition;

Mind. = Mindfulness Condition; *SD*=standard deviation; K=1000; H.S.=high school; G.E.D.=

General Educational Development.

\*Attention-Deficit/Hyperactivity Disorder (ADHD) &/or Anxiety

**Table 2**

*Descriptive statistics for outcomes*

Outcome	Mind. Pre-test		Aud. Pre-test		Aud. + S. Pre-test		<i>p</i>
	<i>n</i>	Mean ( <i>SD</i> )	<i>n</i>	Mean ( <i>SD</i> )	<i>n</i>	Mean ( <i>SD</i> )	
Negative Affect CR	86	14.66(5.23)	88	14.60(5.05)	98	14.63(5.00)	0.99
Mindfulness	90	26.07(6.74)	90	27.58 (6.11)	99	27.96(6.54)	0.11
Stress Children	90	11.99(5.70)	90	10.56(4.33)	99	11.08(5.06)	0.16
Anxiety CR	90	41.90(7.03)	90	41.53(8.07)	99	41.53(7.47)	0.93
Depression CR	90	44.85(7.80)	90	43.89(8.01)	99	43.29(8.50)	0.42
GEC	86	58.01(10.85)	85	55.82(10.98)	94	57.09(12.08)	0.45
BRI	86	55.26(11.19)	85	53.34(11.59)	94	54.22(12.49)	0.57
Anxiety PR	87	47.90 (9.09)	84	45.24(9.82)	92	46.69(10.02)	0.20

Depression PR	87	50.78(10.63)	84	50.02(10.75)	92	50.52(11.86)	0.90
Stress Parent	87	16.02(6.64)	84	15.31(6.80)	92	14.35(7.20)	0.27
Prosociality	87	5.67(2.13)	84	6.06(2.01)	92	5.85(2.26)	0.49
Negative Affect PR	87	7.31(4.08)	84	6.36(3.44)	92	6.25(4.08)	0.14
		<b>Mind. Post-test</b>		<b>Aud. Post-test</b>		<b>Aud. + S. Post-test</b>	
		<i>n</i>		<i>n</i>		<i>n</i>	
		Mean ( <i>SD</i> )		Mean ( <i>SD</i> )		Mean ( <i>SD</i> )	
Negative Affect CR	77	13.56(5.32)	77	13.47(4.95)	86	13.98(5.00)	
Mindfulness	78	27.24(5.56)	77	27.91(7.34)	80	28.42(6.82)	
Stress Children	77	10.39(5.40)	77	10.38(5.28)	82	10.68(5.67)	
Anxiety CR	77	40.75(7.09)	77	40.99(9.11)	82	40.86(8.42)	
Depression CR	77	43.51(9.08)	77	43.12(9.24)	82	42.22(9.50)	
GEC	79	56.53(11.39)	77	55.55(12.10)	85	56.04(11.37)	
BRI	79	53.61(11.05)	77	52.13(11.86)	85	52.98(12.07)	
Anxiety PR	79	46.28(8.84)	74	44.44(9.61)	82	46.60(9.88)	
Depression PR	79	49.08(8.79)	74	48.91(10.39)	82	49.38(10.76)	
Stress Parent	79	14.70(6.43)	74	15.01(7.85)	82	13.24(6.99)	
Prosociality	79	5.80 (2.03)	74	6.28(1.98)	82	6.00(1.87)	
Negative Affect PR	79	5.75(3.72)	74	6.51(3.83)	82	5.43(3.86)	

*Note:* Aud. + S. = Audiobook + Scaffolding Condition; Aud. = Audiobook Only Condition; Mind. = Mindfulness Condition; *SD*=standard deviation; CR: Child Report; PR: Parent Report; Negative Affect CR: Child Negative Affect; Mindfulness: Child and Adolescent Mindfulness Measure (CAMM); Stress Children: Perceived Stress Scale for Children; Anxiety CR: RCADS – Anxiety Scale; Depression CR: RCADS – Depression Scale; GEC: BRIEF Global Executive Composite; BRI: BRIEF Behavioral Regulation Index; Anxiety PR: RCADS – Anxiety Scale; Depression PR: RCADS – Depression Scale; Stress Parent: Perceived Stress Scale; Prosociality: Parent Report Prosociality, Negative Affect PR: Parent Report of Child Negative Affect. *p*-values are derived from ANOVA baseline score comparisons.

**Table 3***Pre-post effect sizes and ANOVA results*


Outcomes					Mind. Vs Aud.		Mind. Vs Aud. + S.		
	$d_{MIND}$	$d_{AUD}$	$d_{AUD+S}$	$d_{diff}$ <i>Mind</i> vs <i>Aud.</i>	ANOVA P value	ANOVA FDR- corrected P value	$d_{diff}$ <i>Mind</i> vs <i>Aud. +</i> <i>S.</i>	ANOV A P value	ANOV A FDR- corrected P value
Negative Affect CR	-0.21	-0.23	-0.13	0.02	0.98	0.99	-0.08	0.57	0.93
Mindfulness	0.19	0.05	0.07	0.14	0.36	0.62	0.12	0.23	0.70
Stress Children	-0.29	-0.04	-0.07	-0.25	0.075	0.45	-0.22	0.18	0.70
Anxiety CR	-0.16	-0.06	-0.08	-0.10	0.36	0.62	-0.08	0.73	0.93
Depression CR	-0.16	-0.09	-0.12	-0.07	0.57	0.86	-0.04	0.78	0.93
GEC	-0.13	-0.02	-0.09	-0.11	0.99	0.99	-0.04	0.87	0.93
BRI	-0.15	-0.10	-0.10	-0.05	0.65	0.86	-0.05	0.93	0.93

Outcomes	Mind. Vs Aud.					Mind. Vs Aud. + S.			
	$d_{MIND}$	$d_{AUD}$	$d_{AUD+S}$	$d_{diff}$ <i>Mind</i> <i>vs Aud.</i>	ANOVA P value	ANOVA FDR- corrected P value	$d_{diff}$ <i>Mind</i> <i>vs</i> <i>Aud. +</i> <i>S.</i>	ANOVA P value	ANOVA FDR- corrected P value
Anxiety PR	-0.18	-0.08	-0.01	-0.10	0.30	0.62	-0.17	0.23	0.70
Depression PR	-0.17	-0.11	-0.10	-0.06	0.89	0.99	-0.07	0.82	0.93
Stress Parent	-0.20	-0.04	-0.16	-0.16	0.25	0.62	-0.04	0.91	0.93
Prosociality	0.06	0.11	0.07	-0.05	0.23	0.62	-0.01	0.60	0.93
Negative Affect PR	-0.39	0.04	-0.21	-0.43	0.004**	0.048*	-0.18	0.11	0.70

\*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$


*Note:* Aud. + S. = Audiobook + Scaffolding Condition; Aud. = Audiobook Only Condition; Mind. = Mindfulness Condition;  $d$ : Cohen's  $d$  effect size; CR: Child Report; PR: Parent Report; Negative Affect CR: Child Negative Affect; Mindfulness: Child and Adolescent Mindfulness Measure (CAMM); Stress Children: Perceived Stress Scale for Children; Anxiety CR: RCADS – Anxiety

Scale; Depression CR: RCADS – Depression Scale; GEC: BRIEF Global Executive Composite; BRI: BRIEF Behavioral Regulation Index; Anxiety PR: RCADS – Anxiety Scale; Depression PR: RCADS –Depression Scale; Stress Parent: Perceived Stress Scale; Prosociality: Parent Report Prosociality, Negative Affect PR: Parent Report of Child Negative Affect.

**Figure 1***CONSORT diagram*

**Figure 2**


*Treatment effects for all groups and all scales*



*Note:* Blue is Mindfulness, Green is Audiobook-scaffolded, Red is Audiobook-only. Error bars reflect standard errors. TimePoint 0 is Pre-test, TimePoint 1 is Post-test. Sample size is 279 individuals. CR: Child Report; PR: Parent Report; Negative Affect CR: Child Negative Affect; Mindfulness: Child and Adolescent Mindfulness Measure (CAMM); Stress Children: Perceived Stress Scale for Children; Anxiety CR: RCADS – Anxiety Scale; Depression CR: RCADS – Depression Scale; GEC: BRIEF Global Executive Composite; BRI: BRIEF Behavioral Regulation Index; Anxiety PR: RCADS – Anxiety Scale; Depression PR: RCADS – Depression Scale; Stress Parent: Perceived Stress Scale; Prosociality: Parent Report Prosociality, Negative Affect PR: Parent Report of Child Negative Affect.

**Figure 3**

*Change in parent-reported child negative affect with median split by days practiced*



\* $p < 0.05$ , \*\* $p < 0.01$

LowPracticeMindful is the group of mindfulness participants at or below the median of days practiced (29). HighPracticeMindful is the group above the median. Error bars reflect standard errors. Sample sizes are 90 for Audiobook-only, 99 for Audiobook-scaffolded, 37 for HighPracticeMindful, 35 for LowPracticeMindful, and 18 mindfulness individuals were missing data for days practiced.

**Supplementary Materials – Table of Contents*****Measure validation: CFA******Fig. S1******Fig. S2******Fig. S3******Fig. S4******Table S1******Table S2******Table S3******Table S4******Table S5******Table S6******References*****Measure validation: CFA**

As we constructed our own measure of negative affect, we conducted a post-hoc confirmatory factor analysis in *lavaan* to identify possible limitations in the measure (Rosseel, 2012). We examined the internal validity of the constructed scale of child negative affect, using responses at baseline. The Negative Affect scale, derived from two proposed factors on thirteen affect items, was upheld. The seven negative affect items had comparable loadings, and the fit was reasonable (CFI=0.921, RMSEA=0.078) (Hu & Bentler, 1999).

Fig. S1

*Distribution of mindfulness days practiced, collected from Inner Explorer*





Fig. S2


*Distribution of mindfulness practices completed, reported by parents*



"1-10": 1 to 10 practices, "11-20": 11 to 20 days practices, etc. "No response" refers to parents who didn't report how many days practiced. Sample sizes are 2,2,17,20,34,15 respectively.

Fig. S3


*Correspondence between Parent Report and objective Inner Explorer metric, dosage*



On the x axis, at the end of the study, parents were asked to report how many practices their children completed. On the y axis, we collected days practiced from Inner Explorer (where >1 click on a mindfulness practice during a day counts as a day practiced).

Fig. S4

*Change in negative affect, by parent-reported number of practices*



\* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$

“1-10”: 1 to 10 completed practices, “11-20”: 11 to 20 completed practices, etc. “No response” refers to parents who didn’t report how many days practiced. Error bars reflect standard errors. Sample sizes are 2,2,17,20,34,90,99,15 respectively.

Table S1

*Cross-measure correlations at baseline*

	Negative Affect CR	Mindfulness	Stress Children	Internalizing CR	Anxiety CR	Depression CR	GEC	BRI	Anxiety PR	Depression PR	Stress Parent	Prosociality
Negative Affect CR												
Mindfulness	-0.34****											
Stress Children	0.44****	-0.36****										
Internalizing CR	0.49****	-0.55****	0.55****									
Anxiety CR	0.45****	-0.53****	0.47****	0.92****								
Depression CR	0.43****	-0.43****	0.50****	0.83****	0.56****							
GEC	0.15*	-0.01	0.21***	0.17**	0.12	0.23***						

	Negative Affect CR	Mindfulness	Stress Children	Internalizing CR	Anxiety CR	Depression CR	GEC	BRI	Anxiety PR	Depression PR	Stress Parent	Prosociality
BRI	0.15*	0.03	0.17**	0.13*	0.08	0.19**	0.90*** *					
Anxiety PR	0.14*	0.01	0.14*	0.17**	0.19**	0.13*	0.34*** *	0.40*** *				
Depression PR	0.11	0.04	0.12	0.06	0.02	0.13*	0.54****	0.52****	0.55*** *			
Stress Parent	0.00	0.08	0.20**	0.09	0.04	0.14*	0.37****	0.37****	0.26*** *	0.41****		
Prosociality	-0.03	-0.06	-0.04	0.02	0.09	-0.10	-0.44****	-0.41****	-0.03	-0.27****	-0.17**	
Negative Affect PR	0.13*	0.12*	0.16**	-0.04	-0.04	0.00	0.52****	0.55****	0.42*** *	0.55****	0.43*** *	-0.36****

\*\*\*\* $p < .0001$ , \*\*\* $p < .001$ , \*\* $p < .01$ , \* $p < .05$

CR: Child Report; PR: Parent Report; Mindfulness: Child and Adolescent Mindfulness Measure (CAMM); Stress Children: Perceived Stress Scale for Children; Anxiety CR: RCADS – Anxiety Scale; Depression CR: RCADS – Depression Scale; GEC: BRIEF Global Executive Composite; BRI: BRIEF Behavioral Regulation Index; Anxiety PR: RCADS – Anxiety Scale; Depression PR: RCADS – Depression Scale; Stress Parent: Perceived Stress Scale; Prosociality: Parent Report Prosociality

Correlations were conducted on the full sample of 279 participants.

Table S2

*ANOVA Results – Effects of Time*

Outcomes	P Value Time	FDR-corrected P Value Time
Negative Affect CR	0.00600**	0.0180*
Mindfulness	0.305	0.305
Stress Children	0.0450*	0.0771
Anxiety CR	0.144	0.157
Depression CR	0.101	0.121
GEC	0.00300**	0.0120*
BRI	0.000417***	0.00500**
Anxiety PR	0.0640	0.0960
Depression PR	0.0110*	0.0220*
Stress Parent	0.00900**	0.0216*
Prosociality	0.0950	0.121
Negative Affect PR	0.000934***	0.00560**

\*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$

*Note:* CR: Child Report; PR: Parent Report; Negative Affect CR: Child Negative Affect; Mindfulness: Child and Adolescent Mindfulness Measure (CAMM); Stress Children: Perceived Stress Scale for Children; Anxiety CR: RCADS – Anxiety Scale; Depression CR: RCADS – Depression Scale; GEC: BRIEF Global Executive Composite; BRI: BRIEF Behavioral Regulation Index; Anxiety PR: RCADS – Anxiety Scale; Depression PR: RCADS – Depression Scale; Stress Parent: Perceived Stress Scale; Prosociality: Parent Report Prosociality, Negative Affect PR: Parent Report of Child Negative Affect

Table S3

*Treatment effects – main ANCOVA results*

Outcomes	Mind vs Aud.			Mind vs Aud. + S.		
	Group B Aud.	P Aud.	FDR P Aud.	Group B Aud.+S.	P Aud.+S.	FDR P Aud.+S.
Negative Affect CR	-0.012	0.987	0.987	0.532	0.449	0.963
Mindfulness	-0.377	0.696	0.987	-0.282	0.767	0.963
Stress Children	0.815	0.278	0.987	0.791	0.271	0.963
Anxiety CR	0.608	0.516	0.987	0.339	0.712	0.963
Depression CR	0.364	0.737	0.987	-0.062	0.953	0.963
GEC	-0.034	0.971	0.987	0.06	0.949	0.963
BRI	-0.548	0.568	0.987	-0.044	0.963	0.963
Anxiety PR	0.226	0.832	0.987	0.83	0.410	0.963
Depression PR	0.289	0.817	0.987	0.611	0.607	0.963
Stress Parent	0.627	0.452	0.987	-0.382	0.637	0.963

Outcomes	Mind vs Aud.			Mind vs Aud. + S.		
	Group B Aud.	P Aud.	FDR P Aud.	Group B Aud.+S.	P Aud.+S.	FDR P Aud.+S.
Prosociality	0.367	0.149	0.894	0.113	0.651	0.963
Negative Affect PR	1.325	0.011*	0.132	0.421	0.399	0.963

\*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$

*Note:* Aud. + S. = Audiobook + Scaffolding Condition; Aud. = Audiobook Only Condition; Mind. = Mindfulness Condition; CR: Child Report; PR: Parent Report; Negative Affect CR: Child Negative Affect; Mindfulness: Child and Adolescent Mindfulness Measure (CAMM); Stress Children: Perceived Stress Scale for Children; Anxiety CR: RCADS – Anxiety Scale; Depression CR: RCADS – Depression Scale; GEC: BRIEF Global Executive Composite; BRI: BRIEF Behavioral Regulation Index; Anxiety PR: RCADS – Anxiety Scale; Depression PR: RCADS – Depression Scale; Stress Parent: Perceived Stress Scale; Prosociality: Parent Report Prosociality, Negative Affect PR: Parent Report of Child Negative Affect

Table S4

*Treatment effects with demographics covarying in ANCOVA models.*

Outcomes	Mind vs Aud.			Mind vs Aud. + S.		
	Group B Aud.	P Aud.	FDR P Aud.	Group B Aud. + S.	P Aud. + S.	FDR P Aud. + S.
Negative Affect CR	-0.232	0.749	0.983	0.423	0.548	0.957
Mindfulness	-0.092	0.925	0.983	-0.052	0.957	0.957
Stress Children	0.874	0.246	0.983	0.941	0.201	0.957
Anxiety CR	0.301	0.749	0.983	0.081	0.93	0.957
Depression CR	0.05	0.964	0.983	-0.268	0.804	0.957
GEC	-0.02	0.983	0.983	0.06	0.948	0.957
BRI	-0.573	0.56	0.983	-0.096	0.919	0.957
Anxiety PR	0.274	0.800	0.983	0.74	0.469	0.957
Depression PR	0.377	0.764	0.983	0.573	0.632	0.957
Stress Parent	0.618	0.467	0.983	-0.361	0.663	0.957
Prosociality	0.281	0.263	0.983	0.03	0.903	0.957
Negative Affect PR	1.44	0.006**	0.072	0.496	0.326	0.957

\*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*  $p < 0.001$

*Note:* Aud. + S. = Audiobook + Scaffolding Condition; Aud. = Audiobook Only Condition; Mind. = Mindfulness Condition; CR: Child Report; PR: Parent Report; Negative Affect CR: Child Negative Affect; Mindfulness: Child and Adolescent Mindfulness Measure (CAMM); Stress Children: Perceived Stress

Scale for Children; Anxiety CR: RCADS – Anxiety Scale; Depression CR: RCADS – Depression Scale; GEC: BRIEF Global Executive Composite; BRI: BRIEF Behavioral Regulation Index; Anxiety PR: RCADS – Anxiety Scale; Depression PR: RCADS – Depression Scale; Stress Parent: Perceived Stress Scale; Prosociality: Parent Report Prosociality, Negative Affect PR: Parent Report of Child Negative Affect

Table S5

*Treatment effects with moderation by demographics in ANCOVA, for child perceived stress*

Moderators	Group X Baseline B Aud.	P Aud.	FDR P Aud.	Group X Baseline B Aud.+S.	P Aud.+S.	FDR P Aud.+S.
Child.Gender	0.919	0.421	0.485	1.247	0.249	0.383
Child.Mental.Health.Dx	0.93	0.288	0.403	0.458	0.588	0.706
WhiteOrNot	1.608	0.256	0.403	1.745	0.206	0.383
IncomeGreater	2.372	0.106	0.403	2.711	0.067	0.383
Educated	0.762	0.485	0.485	0.089	0.933	0.933
Older	1.258	0.228	0.403	1.184	0.255	0.383
SpanishOrNot	1.08	0.158	0.403	0.924	0.223	0.357

Demographic variables are binarized. Child Gender: Male or Female (reported by parent). Child Mental Health Dx.: whether or not a child has a diagnosis of ADHD or anxiety. WhiteOrNot: White or non-white. IncomeGreater: household income greater than 80k. Educated: Maternal Education consisting of masters or above. Older: child age greater than median age (9 years 5 months), SpanishOrNot: parent language spoken, Spanish (1) or English (0).

Table S6

*Treatment effects with moderation by demographics in ANCOVA, for child anxiety*

Outcomes	Group X Baseline B Aud.	P Aud.	FDR P Aud.	Group X Baseline B Aud.+S.	P Aud.+S.	FDR P Aud.+S.
Child.Gender	1.336	0.349	0.611	1.462	0.281	0.550
Child.Mental.Health.Dx	0.681	0.534	0.748	0.922	0.393	0.550
WhiteOrNot	2.854	0.108	0.410	1.687	0.336	0.550
IncomeGreater	0.178	0.924	0.924	2.067	0.267	0.550
Educated	0.134	0.92	0.924	0.523	0.688	0.785
Older	2.056	0.117	0.410	1.798	0.172	0.550
SpanishOrNot	1.037	0.285	0.611	0.263	0.785	0.785

Demographic variables are binarized. Child Gender: Male or Female (reported by parent). Child Mental Health Dx.: whether or not a child has a diagnosis of ADHD or anxiety. WhiteOrNot: White or non-white. IncomeGreater: household income greater than 80k. Educated: Maternal Education consisting of masters or above. Older: child age greater than median age (9 years 5 months). SpanishOrNot: parent language spoken, Spanish (1) or English (0).

## References

Hu, L. T., & Bentler, P. M. (1999). Cutoff criteria for fit indexes in covariance structure analysis:

Conventional criteria versus new alternatives. *Structural equation modeling: a multidisciplinary journal*, 6(1), 1-55.

Rosseel, Y. (2012). lavaan: An R package for structural equation modeling. *Journal of Statistical*

*Software*, 48(2), 1-3.