

Major Lifetime Discrimination Exposure Accelerates Brain Microstructural Aging: Diffusion MRI Evidence From MIDUS

Ajay Kumar Nair, PhD, Nagesh Adluru, PhD, Anna J. Finley, PhD, Lauren K. Gresham, BA, Sarah E. Skinner, BS, Steven R. Kecskemeti, PhD, Andrew L. Alexander, PhD, Richard J. Davidson, PhD, David R. Williams, PhD, Carol D. Ryff, PhD, and Stacey M. Schaefer, PhD

Objective: Experiencing discrimination is associated with faster biological aging, as reflected in telomere shortening and DNA methylation. However, the impact of discrimination on brain aging processes remains unclear. Here, we tested whether individuals who reported at least one major lifetime discrimination event would exhibit steeper age-related associations in microstructural metrics within whole-brain white matter and the hippocampus, consistent with accelerated brain microstructural aging, compared with those with no such experiences.

Methods: We analyzed multi-shell diffusion-weighted MRI data from the Midlife in the United States (MIDUS) cohort ($n = 147$, mean age = 65 years, range: 48 to 95 years) to assess brain micro-

structure using complementary statistical and biophysical diffusion models. Diffusion kurtosis imaging representation was used to derive diffusion tensor imaging (DTI) and white matter tract integrity (WMTI) measures. Additional microstructural health indices were derived using the neurite orientation dispersion and density imaging (NODDI) model. Permutation analyses of linear models were run within the whole-brain white matter and bilateral hippocampi, adjusting for sex, race, and education.

Results: Participants who reported at least one major discriminatory experience during their lifetime exhibited accelerated age-associated changes in white matter microstructural measures, including higher mean and radial diffusivities, extra-axonal radial diffusivity, and free water fraction compared with those with no such experiences.

Conclusions: These converging findings from complementary measures of brain microstructure suggest that major discrimination experiences may contribute to accelerated brain microstructural aging.

Key Words: major lifetime discrimination, aging, diffusion weighted imaging, brain microstructure, MIDUS

Abbreviations: BIPOC = Black, Indigenous, and People of Color, CSF = cerebrospinal fluid, DKI = diffusion kurtosis imaging, DTI = diffusion tensor imaging, DWI = diffusion weighted imaging, FA = fractional anisotropy, FOV = field of view, FWE = family-wise error, MD = mean diffusivity, MIDUS = Midlife in the United States, MK = mean kurtosis, MRI = magnetic resonance imaging, NODDI = neurite orientation dispersion and density imaging, PALM = permutation analyses of linear models, PET = positron emission tomography, RD = radial diffusivity, RK = radial kurtosis, TE = echo time, TR = repetition time, WMTI = white matter tract integrity

(*Biopsychosoc Sci Med* 2026;88:297–309)

INTRODUCTION

Discrimination is a social determinant of health with profound negative impacts, including increased risk for cardiovascular disease, depression, psychological dis-

Received for publication August 4, 2025; accepted December 15, 2025.

From the Institute on Aging, University of Wisconsin-Madison, Madison, Wisconsin (Nair, Adluru, Finley, Gresham, Skinner, Ryff, Schaefer); Centre for Brain and Mind, National Institute of Mental Health and Neurosciences, Bengaluru, India (Nair); Waisman Center, University of Wisconsin-Madison, Madison, Wisconsin (Adluru, Kecskemeti, Alexander); Department of Radiology, University of Wisconsin-Madison, Madison, Wisconsin (Adluru); Department of Psychology, North Dakota State University, Fargo, North Dakota (Finley); Department of Medical Physics, University of Wisconsin-Madison, Madison, Wisconsin (Alexander); Department of Psychiatry, University of Wisconsin-Madison, Madison, Wisconsin (Alexander, Davidson); Department of Psychology, University of Wisconsin-Madison, Madison, Wisconsin (Davidson, Ryff); Center for Healthy Minds, University of Wisconsin-Madison, Madison, Wisconsin (Davidson); and Department of Social and Behavioral Sciences, Harvard T. H. Chan School of Public Health, Boston, Massachusetts (Williams).

Article Editor: Suzanne C. Segerstrom

Address correspondence to Stacey M. Schaefer, PhD, MIDUS Neuro, 901 University Bay Dr Madison, WI 53705. E-mail: stacey.schaefer@wisc.edu

Supplemental Digital Content is available for this article. Direct URL citations are provided in the HTML and PDF versions of this article on the journal's website, journals.lww.com/bsam.

Copyright © 2026 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the Society for Biopsychosocial Science and Medicine. This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

DOI: 10.1097/PSY.0000000000001467

stress, suicidal ideation, and mortality.¹ Experiencing discrimination is stressful. The biological weathering hypothesis suggests that exposure to psychosocial stressors such as discrimination can accelerate the aging process, increasing vulnerability to disease and early mortality.² For example, experiencing discrimination is positively associated with both heightened threat appraisals and greater negative affective reactivity to stress that mediate worse physical and mental health outcomes 20 years later.³ However, whether experiences of discrimination can prematurely age the brain remains to be studied.

Major discriminatory events—such as being fired from a job, hassled by the police, or denied a promotion due to an individual characteristic such as age, sex, race, ethnicity, disability—are widespread in the United States. At least one-third of the population reports experiencing major discrimination during their lifetime.^{4,5} Discrimination has detrimental consequences that are particularly relevant to aging. Age is one of the strongest risk factors for many neurodegenerative conditions, yet the pace of biological aging may differ from chronological age.⁶ Major discrimination has been associated with greater biological aging, as indicated by shorter telomere length.⁷ Individuals who reported more frequent or multiple forms of discrimination showed older biological ages and a faster pace of aging, based on blood DNA methylation markers.⁸ Other adverse outcomes associated with discrimination, and relevant to aging, include smaller hippocampal volume⁹ and poorer cognition, including worse episodic memory¹⁰ and reduced executive function.¹¹

Age is also strongly associated with declining brain health among older individuals, with reductions in volumes of most brain structures, especially the hippocampus, accelerating from the fifties onwards.^{12,13} Even before volumetric changes or neurodegenerative conditions become detectable in the hippocampus and other brain regions, subtle microstructural changes can be identified using diffusion-weighted magnetic resonance imaging (DWI).^{14,15} However, the association between major discrimination and brain microstructure in the context of aging has not yet been examined.

By leveraging the diffusion properties of water molecules, DWI can be used to derive measures that are sensitive to cellular microstructure such as changes caused by edema or necrosis.¹⁶ Statistical models based on signal representations, such as diffusion tensor imaging (DTI) and diffusion kurtosis imaging (DKI), can be applied to each voxel to obtain measures dependent on the normal distribution of water displacement, including: (1) fractional anisotropy (FA), reflecting the extent of diffusivity in a preferential direction such as along axonal fibers, (2), mean diffusivity (MD), the average diffusivity in all directions, and (3) radial diffusivity (RD), the diffusivity perpendicular to axonal fibers, or complementary measures such as (4) mean kurtosis (MK) and (5) radial kurtosis (RK) that indicate deviations from the normal displacement distribution.

Advanced biophysical models, such as neurite orientation dispersion and density imaging (NODDI) and

white matter tract integrity (WMTI), represent the underlying tissue microstructure by separating each voxel into compartments with restricted and hindered diffusion of water molecules representing diffusion of water molecules within axons or in extracellular space.^{17,18} For example, the NODDI model estimates (1) neurite density index, representing density of axons and dendrites (collectively called neurites), (2) orientation dispersion index, which indicates if these neurites are tightly packed or spread out, and (3) proportion of cerebrospinal fluid (CSF) estimated as the free water fraction.¹⁸ The WMTI model estimates measures such as: (1) axonal water fraction, related to axonal density, (2) intra-axonal diffusivity, capturing restricted diffusion within myelinated axons, and (3) extra-axonal diffusivities, including hindered diffusion alongside the axon (typically higher than intra-axonal diffusivity) and perpendicular to the axon (typically lower diffusivity than along the axon).¹⁷

Taken together, the above measures provide complementary and overlapping information and are sensitive to brain microstructural changes in myelination and axonal density, across the lifespan, from early brain development¹⁹ to aging and neurodegenerative conditions.^{20,21} A comprehensive assessment using multiple microstructural metrics can reveal distinct patterns of findings that, when considered together, can help infer specific biological processes that underlie associations with the predictor of interest (e.g., Nair et al.²²), such as experiencing a major lifetime discriminatory event.

Using complementary parameters from statistical models (DTI and DKI), and biophysical models (WMTI and NODDI), we investigated brain microstructural features associated with age across a wide age range and examined how experiencing at least one major discriminatory event compared with no experiences of major discrimination moderated age associations with DWI measures. Brain regions showing converging findings across multiple DWI models and measures robustly reflects moderation of age-related changes with a discrimination experience. We specifically examined microstructural features in two regions of interest: (1) the whole-brain white matter, and (2) the hippocampus, given its previous associations suggesting vulnerability to both age and discrimination experiences. We hypothesized that experiencing at least one major discrimination event would be associated with steeper age-related associations in microstructural metrics within whole-brain white matter and the hippocampus, consistent with accelerated brain microstructural aging.

METHODS

Study Overview

We used data from the Midlife in the United States (MIDUS) longitudinal study, which recruited baseline participants aged 25 to 74 years in 1995/96 and followed them at multiple timepoints. An oversample of Black/African American participants from Milwaukee, Wisconsin was added during the second follow-up (MIDUS2; 2004 to 2009). MIDUS seeks to understand the interplay

of sociodemographic, psychosocial, and neurobiological determinants of health and disease among aging adults. MIDUS data are publicly available for use by scientists around the world (<https://midus.wisc.edu/>). During the third follow-up of the MIDUS Core sample, multi-shell DWI data were acquired for the first time by the MIDUS Neuroscience Project (MIDUS3; 2017-2022). All participants traveled to Madison, Wisconsin and were scanned on the same MRI scanner. Only MIDUS3 data were used for the cross-sectional analyses reported in the present work. Approval was obtained from relevant institutional review boards. Participants were briefed on study procedures, screened for MRI compatibility, and provided informed consent before data collection.

Participant Characteristics

Participants ($n = 147$) were between the ages of 48 to 95 (mean = 65.03, median = 64, SD = 9.35) years. There were 87 females (59%) and 40 Black, Indigenous, and People of Color (BIPOC, 27.2%) in the sample. Among the BIPOC group, most participants were Black (80%) with very few reporting other racial minorities (2 Native American/Alaska Native, 2 Asian American, and 4 Other). There were 64 participants with a college degree (43.5%), 45 with some college education (30.6%), and 38 with a high school education or less (25.9%).

Major Experiences of Discrimination Assessment

The Major Experiences of Discrimination Scale was administered at MIDUS3.^{23,24} The scale assesses major discriminatory experiences across multiple settings, including academics (discouraged from continuing education), employment (denied promotion), financial services (prevented from renting or buying a home), health care (denied or provided inferior medical care), and social hostility (hassled by the police). Table S1, Supplemental Digital Content, <http://links.lww.com/PSYMED/B161>, presents the full list of 11 questions included in the scale. Participants responded to each item by providing the frequency (number of times) they experienced this type of discrimination over their lifetime, and the item responses were summed to create a total score. The scale has been validated and used widely in prior research.^{25–27} In the current sample, scores ranged from 0 to 9, with a median of 0 and a mean score of 1.177. Participants were categorized into 2 groups: those who reported no major experiences of discrimination (no discrimination group; score = 0), and those who reported experiencing at least one major discriminatory event (major discrimination group; score ≥ 1). Table S2, Supplemental Digital Content, <http://links.lww.com/PSYMED/B161>, presents sample characteristics stratified by major discrimination group status. Individuals identifying as BIPOC were significantly more likely to report having experienced major discrimination compared with White participants. No other group differences reached statistical significance. Within the segment of the sample ($n = 64$; 43.5%) that reported experiencing at least one major discriminatory

event, the median score was 2 and the mean was 2.7. Individual responses to the major lifetime discrimination questions spanned several categories (17 hassled by the police, 17 fired, 13 participants reported being discouraged from higher education, 12 denied or provided inferior service, 9 denied a bank loan, 8 prevented from renting/buying a home, 5 denied or provided inferior medical care, 3 denied a scholarship and 2 prevented from remaining in a neighborhood). Available attributions for overall discrimination experience, not specific to major discrimination, were diverse in both BIPOC and White participants with some overlap. Table S3, Supplemental Digital Content, <http://links.lww.com/PSYMED/B161>, presents attributions of overall discrimination experience stratified by race.

DWI Data Acquisition

A multi-shell spin-echo, echo-planar imaging sequence was used to collect DWI data on a 3 Tesla GE MR750 scanner with a Nova 32-channel head coil. Three shells of different diffusion-weighting strengths were acquired at b-values of 500, 800, and 2000 s/mm² with 9, 18, and 36 directions, respectively. There were 6 reference scans without any diffusion encoding ($b = 0$ s/mm²). Other parameters included: repetition time/echo time (TR/TE) = 7000/91 ms; field of view (FOV) = 256 mm; 75 slices; voxel resolution = 2×2×2 mm³.

DWI Analysis

Figure S1 (Supplemental Digital Content, <http://links.lww.com/PSYMED/B161>) provides an overview of the multi-model analysis approach used in the study.

Pre-Processing and Estimation of Brain Microstructure Metrics

The DESIGNER guidelines were used for pre-processing the data.²⁸ The pre-processing steps included the removal or mitigation of artifacts such as noise, Gibb's ringing, distortion because of eddy currents, and B1 and "Rician" bias correction using tools in FSL v6.0,²⁹ ANTS,³⁰ and MRtrix3.³¹ After weighted least-squares optimization for voxel-wise estimation of the diffusion kurtosis tensor and the diffusion tensor³² in a joint estimation based on the Taylor series decomposition of the signal, image maps of the DKI and DTI metrics were generated. The WMTI model parameters were computed from the estimates of the kurtosis tensor. Gaussian smoothing of 2 voxels (full width half maximum of 4.7096) was uniformly used for DTI and NODDI maps. To address spatial variation in kurtosis estimates, spatially adaptive smoothing was applied to the DKI and WMTI measures²² as follows. A mask was created using a threshold MK < 0.3 and was smoothed to effectively obtain a "weighting" map. Then each of the DKI and WMTI metric maps was smoothed and divided by the smoothed weighting mask resulting in smoothed metrics that downweigh the influence of the low-MK/noisy DKI estimates (also known in the literature as "black holes").^{33–35} This approach was used as a refinement to

the typical solution of isotropic smoothing of the DKI metrics, inspired by the T-SPOON method,³⁶ which aims to reduce the influence of misregistration errors in voxel-based analysis.

In addition, the multi-tissue NODDI model³⁷ was fit to the DWI data to derive voxel-wise estimates of corresponding measures. The multi-tissue response functions used in estimating the NODDI parameters help account for the different tissue types (white matter, gray matter, and CSF) in mitigating biases in the derived measures.³⁷ Quality control was performed visually to ensure proper orientations of the images, brain masking, and to ensure the quality of the estimated parameter maps looks typical and appropriate without artifacts such as black holes. All parameter maps were warped to a study-specific population template that was estimated using individual subject FA and MD maps from all the participants.³⁸ Parameter maps from all participants were grouped measure-wise and merged for statistical analyses.

Statistical Analyses and Visualization

Permutation analyses of linear models (PALM) examined differences in DWI microstructure metrics by major lifetime discrimination experience.³⁹ For primary analyses, models examined the moderating influence of discrimination group status (major lifetime discrimination group vs. no major lifetime discrimination group) on age-associated changes in microstructure (ie, included a discrimination group \times age interaction term) adjusting for sociodemographic covariates: sex (male, female), race (White, BIPOC), educational attainment level (high school or less, some college, college degree). The analyses were restricted to voxels in two regions of interest (ROIs) using the following approach: (1) white matter skeleton in the whole-brain mask provided with FSL⁴⁰ was warped to the study population template space and binarized. Note that the skeleton was not applied within the TBSS framework⁴¹ but was instead used purely as a voxel mask for the analyses. (2) Bilateral hippocampal regions from the Harvard-Oxford subcortical atlas⁴² were warped to the study population template space. Analyses were run using analytical tail acceleration and 500 permutations. Joint inference was carried out for each set of models (DTI, DKI, WMTI, and NODDI) using non-parametric combination, along with simultaneous inference for each individual metric. Threshold-free cluster enhancement and family-wise error (FWE) correction for multiple comparisons were used to control for false positives at an $\alpha < 0.05$, and corresponding statistical brain maps were generated. The XTRACT HCP Probabilistic Tract Atlas from FSL was used to identify regions with significant findings.

To visualize significant associations between microstructure metrics and variables of interest, the mean value of each outcome metric was computed from all significant voxels within each ROI for each participant. These participant-level averages were then used as summary outcome measures for visualization. Corresponding linear models were run, and partial residuals were plotted

for significant associations. Model coefficients presented in the results section are for descriptive purposes as these models utilize data extracted from voxels found significant after statistical testing. Influential outliers were defined as those exceeding a threshold of 5% of the F-distribution of Cook's distance and were removed. All results were consistent with and without outlier removal. Visualizations were carried out using R statistical software v4.4.0.⁴³ To illustrate spatial convergence across models, conjunction maps were generated to visualize the overlap of significant age \times discrimination group interaction effects across DTI, WMTI, and NODDI metrics.

RESULTS

Major lifetime discrimination group status significantly moderated the associations between age and microstructural measures derived from DTI, WMTI, and NODDI, in both the whole-brain white matter skeleton and the hippocampus, after adjusting for the sociodemographic covariates (sex, race, and educational attainment). No significant associations were observed for the DKI parameters. Unless otherwise specified, findings reported were consistent with and without inclusion of sociodemographic covariates in the models. Table S4, Supplemental Digital Content, <http://links.lww.com/PSYMED/B161>, summarizes the significant findings across the tested models. All significant models exhibited a sign/directionality consistent with an age-moderation effect suggestive of accelerated microstructural aging. None of the models with inverse contrasts were statistically significant. Note that counts or frequency of major lifetime discrimination experience was not significantly associated with age in this sample ($r = -0.07$, $P = .40$; see Figure S3, Supplemental Digital Content, <http://links.lww.com/PSYMED/B161>, but age was significantly associated with all white matter metrics reported below (see Table S5, Supplemental Digital Content, <http://links.lww.com/PSYMED/B161>). Considerable overlap was observed in white matter voxels showing significant age \times discrimination group interactions, as shown in the conjunction map across DTI, WMTI, and NODDI models (Figure S2, Supplemental Digital Content, <http://links.lww.com/PSYMED/B161>).

Age-Discrimination Interactions in Whole-Brain White Matter: Statistical Models

Two statistical models were used to examine age-discrimination interactions in the whole-brain white matter skeleton: diffusion tensor imaging (DTI) and diffusion kurtosis imaging (DKI). For DTI, fractional anisotropy, mean diffusivity, and radial diffusivity were tested, and for DKI, mean kurtosis and radial kurtosis were assessed.

Individuals who reported experiencing at least one major discriminatory experience showed higher mean diffusivity values with greater age ($\beta = 0.00362$, $P < .001$, one outlier removed), compared with those with no such experiences. These associations were observed across widespread regions, including the anterior and superior




FIGURE 1. Associations between white matter diffusion tensor metrics and age by major lifetime discrimination group status. Scatter plots visualizing the moderating influence of major lifetime discrimination group status on the voxel-wise relationships between age and (A) mean diffusivity, and (C) radial diffusivity within the whole-brain white matter mask. Data points are color-coded and represented as triangles for those who experienced major discrimination, and circles for those who did not. Each data point represents the mean of all significant voxels for one individual, adjusted for sex, race, and educational attainment. Representative brain slices show voxels with significant relationships (at $\alpha < 0.05$, family-wise error corrected) for (B) mean diffusivity and (D) radial diffusivity. Brain images are in radiologic convention (left hemisphere is shown on the right side). *Color image is available only in online version.*

thalamic radiations, forceps minor, corticospinal tract, inferior fronto-occipital fasciculus, and acoustic and optic radiations (Figure 1A, B).

A similar pattern was observed for radial diffusivity ($\beta = 0.00428$, $P < .001$), with higher values associated with greater age in the anterior and superior thalamic radiations, forceps minor, corticospinal tract, inferior fronto-occipital fasciculus, superior longitudinal fasciculus, dorsal cingulum subsection, and the acoustic radiation (Figure 1C, D). No significant age-discrimination interactions were observed for DKI metrics.

Age-Discrimination Interactions in Whole-Brain White Matter: Biophysical Models

Two biophysical models were used to examine age-discrimination interactions in the whole-brain white matter skeleton: neurite orientation dispersion and density imaging (NODDI) and white matter tract integrity (WMTI). For NODDI, neurite density index, orientation dispersion index, and CSF fraction were tested. For WMTI, axonal water fraction, intra-axonal diffusivity, extra-axonal radial diffusivity, and extra-axonal tortuos-

ity were assessed.

Individuals who reported at least one major discriminatory experience showed steeper associations between age and the NODDI metric CSF fraction ($\beta = 0.00215$, $P < .001$) compared with those with no experiences of discrimination. These associations were observed in widespread regions including the anterior and superior thalamic radiations, forceps major, corticospinal tract, inferior fronto-occipital fasciculus, superior longitudinal fasciculus, arcuate fasciculus, optic radiation, and the vertical occipital fasciculus (Figure 2A, B).

A similar pattern was observed for the WMTI metric extra-axonal radial diffusivity ($\beta = 0.00348$, $P < .001$, one outlier removed), with higher values associated with greater age in the anterior and superior thalamic radiations, forceps minor, corticospinal tract, dorsal cingulum subsection, and the acoustic and optic radiations (Figure 2C, D).

Although the association with CSF fraction remained consistent, the association with extra-axonal radial diffusivity was not significant when sociodemographic covariates were dropped.




FIGURE 2. Associations of white matter microstructural metrics from biophysical models and age by major lifetime discrimination group status. Scatter plots visualizing the moderating influence of major lifetime discrimination group status on the voxel-wise relationships between age and (A) CSF fraction, and (C) extra-axonal radial diffusivity in the whole-brain white matter mask. Data points are color-coded and represented as triangles for those who experienced discrimination, and circles for those who did not. Each data point represents the mean of all significant voxels for one individual, adjusted for sex, race, and educational attainment after removing influential outliers, if any. There was one influential outlier for the model with extra-axonal radial diffusivity. Results were consistent with ($\beta=0.0046$) and without the outlier ($\beta=0.0034$). Representative brain slices show voxels with significant relationships (at $\alpha < 0.05$, family-wise error corrected) for (B) CSF fraction and (D) extra-axonal radial diffusivity. Brain images are shown in radiologic convention (left hemisphere is shown on the right side). *Color image is available only in online version.*

No significant age-discrimination interactions were observed for WMTI metrics related to axonal water fraction, intra-axonal diffusivity, or extra-axonal tortuosity. Similarly, no significant interactions were found for NODDI metrics related to neurite density or orientation dispersion. However, a trend-level effect was observed for neurite density ($P = .089$), which is further discussed below.

Age-Discrimination Interactions in Hippocampal Microstructure

Analyses within the hippocampal region of interest (ROI), which included both left and right hippocampi, examined age-discrimination interactions using diffusion metrics derived from the statistical models (DTI, DKI) and biophysical models (NODDI, WMTI). For DTI, mean diffusivity, radial diffusivity, and fractional anisotropy were tested. For DKI, mean kurtosis and radial kurtosis were assessed. For NODDI, neurite density index, orientation dispersion index, and CSF fraction were

tested. For WMTI, axonal water fraction, intra-axonal diffusivity, extra-axonal radial diffusivity, and extra-axonal tortuosity were assessed.

Older participants who had experienced major discrimination showed higher extra-axonal radial diffusivity in the right hippocampus ($\beta = 0.00694$, $P < .001$; Figure 3A, B), whereas no significant association with age was found in participants without major discrimination experience.

No significant age-discrimination interactions were observed for other metrics in the hippocampal ROI.

DISCUSSION

The present study tested the hypothesis that experiences of major lifetime discrimination moderate age-related differences in brain health, as indicated by DWI measures of whole-brain white matter and hippocampal microstructure. Supporting this hypothesis, we found that individuals who reported major lifetime discrimination



FIGURE 3. Associations of hippocampal microstructure metrics and age by major lifetime discrimination group status. A, Scatter plot visualizing the moderating influence of major lifetime discrimination group status on the voxel-wise relationships between age and extra-axonal radial diffusivity in the bilateral hippocampal mask, adjusting for sex, race, and educational attainment. Data points are color-coded and represented as triangles for those who experienced discrimination, and circles for those who did not. Each data point represents the mean of all significant voxels for one individual. B, Representative brain slices show voxels with significant relationships (at $\alpha < 0.05$, family-wise error corrected). *Color image is available only in online version.*

(having experienced at least one severe discriminatory event such as being denied a job, a promotion, or a loan due to discrimination) showed steeper relationships between age and microstructural health consistent with accelerated brain aging. These effects were observed across widespread regions of whole-brain white matter and in the hippocampus. Older age is a powerful predictor of adverse brain microstructural changes. However, major lifetime discrimination experience amplified (exacerbated) relationships between age and complementary diffusion parameters from both statistical and biophysical diffusion models in white matter and the hippocampus.

A growing literature describes the adverse effects of discrimination, including accelerated biological aging and increased risk for dementia.^{7,8,44–46} These studies suggest multiple mechanisms, such as higher cardiometabolic risk, increased inflammation and overall allostatic load, and epigenetic dysregulation, by which discrimination may accelerate biological aging. The present study adds to this literature by identifying microstructural differences, suggesting that discrimination experience may accelerate brain aging. We found that participants who experienced at least one major discriminatory experience in their lifetime had greater values of mean and radial diffusivities with age across widespread white matter regions. These regions also showed greater values of CSF fraction and extra-axonal radial diffusivity. As shown by the conjunction analysis in the supplementary material, Supplemental Digital Content, <http://links.lww.com/PSYMED/B161>, most findings across the DWI metrics were in consistent, overlapping regions, suggesting that with greater age, participants who reported experiencing one or more major discriminatory events had higher free water content in these white matter regions than was evident for participants of the same age who had not experienced any such events. Increased brain interstitial water can occur through multiple pathways, including

higher cerebrovascular pathology,⁴⁷ edema,⁴⁸ neuroinflammation,⁴⁹ or white matter fiber shrinkage.^{14,50} Future research should assess which of these pathways mediate the relationship between discrimination and accelerated brain aging.

In a few white matter tracts, such as portions of the forceps minor, significant voxels were overlapping for radial diffusivity and extra-axonal radial diffusivity, but not for the CSF fraction. Higher values of these radial diffusivity measures, one from DTI and the other a more biophysically informed measure from WMTI, indicates greater movement of water perpendicular to the axonal fibers, and is suggestive of demyelination.^{51,52} In addition, participants who experienced major lifetime discrimination had higher values of extra-axonal radial diffusivity in the hippocampus, although no other measures were sensitive to this effect. Interpreting this finding as demyelination is challenging because WMTI measures are less reliable in regions like the hippocampus, which contains white matter bundles but is predominantly gray matter. This is due to water exchange between neurites and extracellular space as well as signal contributions from neuronal somas.⁵³ The hippocampus is also located adjacent to the temporal horn of the lateral ventricles, making it prone to CSF contamination, especially with age-related atrophy. Nevertheless, biophysical models have shown promise in characterizing hippocampal microstructure.^{54–56} Despite these caveats, this result is noteworthy given the important role of the hippocampus in brain health in the context of aging and discrimination due to its role in learning and memory, sensitivity to stress and inflammation, and vulnerability to volume loss with age and neurodegenerative conditions such as Alzheimer disease.^{57,58}

The examination of multiple microstructural imaging models and measures of brain health offers unique opportunities to identify which biological processes may

or may not be involved. For example, increased radial diffusivity from DTI is classically suggestive of demyelination. However, DTI is nonspecific and sensitive to any change that increases perpendicular water motion, including increased extra-axonal space. Biophysical models allow us to test the demyelination/axonal loss hypothesis more directly. A primary mechanism of demyelination or axonal loss would be expected to alter the neurite density index from NODDI. However, we did not find strong evidence that major discrimination experience amplified age-related decline in neurite density (the association showed a trend, $P < .1$). Conversely, we found strong, significant effects in measures of the extra-axonal space, specifically the CSF fraction from NODDI and extra-axonal radial diffusivity from WMTI. This suggests that processes such as demyelination or axonal loss⁵⁹ could be relevant, but are unlikely to be the primary mechanism linking discrimination to brain health. Instead, our findings suggest that the radial diffusivity finding is driven by changes in the extra-axonal space, such as increased interstitial water, which could be related to neuroinflammation, edema, or impaired glymphatic clearance. Larger and/or longitudinal studies will be important to determine whether subtle neurite alterations contribute to discrimination-related brain changes, as would the collection of more direct imaging modalities such as myelin water imaging.⁶⁰ Similarly, major discrimination was not significantly associated with the DKI metrics, despite the significant age-linked associations with NODDI and WMTI metrics. One possible explanation is that NODDI and WMTI derive biologically relevant parameters such as neurite density, orientation dispersion, and free water, by compartmentalizing tissue environments within each voxel into intra- (restricted diffusion) and extra- (hindered diffusion) axonal spaces. Here, restricted diffusion spaces model thermal motion of water that is physically blocked by impermeable or semi-permeable barriers (eg, axonal membranes), whereas hindered diffusion spaces model free water motion that is not confined but is slowed down due to obstacles in the environment (eg, glial cells, convoluted geometry of extracellular spaces).⁶¹ DKI, in contrast, estimates signal deviations from Gaussian diffusion without explicitly modeling tissue compartments and provides markers of overall microstructural complexity.⁶² The absence of DKI effects in our analysis suggests that discrimination-related differences are not generalized alterations in tissue complexity but may reflect more specific microstructural alterations (eg, axonal or myelin alterations) that are more sensitively captured by biophysical models such as NODDI and WMTI.

The white matter tracts where we observed convergent findings also suggest potential impacts of major lifetime discrimination on cognition. The inferior fronto-occipital fasciculus has been linked to various cognitive functions, including perception, context-dependent cognitive control, embodied cognition, and social cognition.^{63,64} The forceps minor has also been associated with social cognition.⁶⁵ The anterior thalamic radiation is implicated in embodied cognition, processing speed, and

set shifting.^{64,66} Taken together, these findings suggest that major discrimination may be associated with accelerated age-related cognitive decline. The MIDUS study is currently following this sample up for the fourth timepoint, and their longitudinal cognitive and imaging data will provide an opportunity to examine the import of these discrimination and age-related microstructural findings for cognitive decline.

Because the timing, context, and frequency of discriminatory experiences can vary greatly, adverse associations of discrimination with health outcomes may also vary by participant background.⁶⁷ For example, although in a multi-ethnic sample discrimination was associated with higher risk for dementia,⁴⁴ a study among only Black Americans did not find any associations with dementia risk,⁶⁸ potentially because in that sample, those with higher discrimination experience had higher education and income, factors that are protective against cognitive impairment and dementia.⁶⁹ Alternatively, a study including only non-Hispanic Black people found that racial discrimination in major life domains was associated with smaller hippocampal volumes, and everyday racial discrimination was associated with faster accumulation of white matter hyperintensities, although discrimination by all causes taken together did not show these associations.⁹

Sources and attributions of discrimination in the United States may vary by sociodemographic group and may differ for daily versus major lifetime discrimination experiences. Prior studies suggest that White individuals often attribute daily discrimination to sex and age, whereas BIPOC individuals more commonly attribute it to race, along with age, sex, and sexuality.^{70,71} As noted earlier, some evidence suggests that both daily and lifetime racial discrimination may be particularly detrimental to brain health, but discrimination regardless of attribution might not show the same effect.⁹ As described in the supplemental material, Supplemental Digital Content, <http://links.lww.com/PSYMED/B161>, discrimination attributions (self-reported broadly for both daily and lifetime discrimination) were highly variable for both White and BIPOC groups in this sample.

Another consideration is that major lifetime experiences of being discouraged from higher education and/or denied a scholarship may directly impact educational attainment, and educational attainment was included as a covariate in our models. Indeed, a subset of our participants (~11%) reported being discouraged from pursuing higher education or denied scholarships. However, most of our findings were consistent with and without adjusting for covariates, and educational attainment was not a significant predictor in any of the white matter models, suggesting that this overlap does not bias our findings. Education remains important for brain health (and overall health and longevity), and needs to be accounted for, especially given the heterogeneity in the sample. Finally, our focus on the presence of any major discriminatory experience, regardless of attribution, is intentional, given the limited number of participants with both imaging and discrimination data, which precludes examining specific

types or sources of discriminatory experience. Given the potential links between discrimination, educational attainment, and other covariates, we interpret adjusted estimates to be robust to potential confounding, but conditional on sociodemographic profile (i.e., holding covariates constant). Future work with larger samples should explore how sociodemographic factors may moderate the effects of discrimination on brain health and how intersectorality may be especially important.

Experiences of discrimination contribute significantly to generalized stress, and are associated with more chronic conditions, greater functional limitations, and faster epigenetic aging.^{8,46,72} One plausible pathway linking these effects is inflammation. Discrimination is associated with higher systemic inflammation and, in turn, with faster memory decline among older adults.⁷³ Markers of systemic inflammation can activate proinflammatory signaling pathways in the brain and lead to accelerated brain aging.^{8,74} In addition, discrimination experiences of any kind may lead to heightened vigilance and threat perception, as suggested by higher resting amygdala activity and higher amygdala connectivity with regions within the salience network.⁷⁵ Prior experiences of discrimination can also prime individuals to anticipate discriminatory behavior even when none exists.⁷⁶ In addition, the presence of concurrent mental health challenges can exacerbate outcomes—for example, individuals who experienced both racial discrimination and depressive symptoms had lower total brain matter and white matter volumes than those who experienced neither.⁷⁷

Future research needs to better understand how discrimination combines with other stressful life experiences to affect accelerated brain aging. Discrimination is only one type of stressor that can adversely affect health, and prior research reveals that experiences of discrimination, like other stressors, are not randomly distributed: socially disadvantaged populations have higher levels of exposure. For example, a study of a probability sample of over 3000 adults in Chicago found that Black/African Americans and US-born Hispanics had higher levels of 3 indicators of discrimination compared with White individuals, and also had higher levels of multiple other stressors [major life events (eg, death of a loved one), financial stress, relationship stress, work stress and neighborhood stress]; each additional domain of stress was associated with worse physical and mental health.⁷⁸ Another priority for future research is to identify how discrimination and other stressors combine, starting early in life, and accumulate over time to affect brain functioning and aging across the life course. The stress acceleration hypothesis indicates that, in response to threatening environments, the brain adapts by expediting neurodevelopment.^{79,80} For example, youth in socioeconomically deprived neighborhoods show reductions in cortical thickness and gray matter volume.^{81,82} Our current understanding is limited regarding how these processes evolve over the life course, and how and when we might optimally intervene to reverse or mitigate them.

A critical challenge for those experiencing major discrimination is that they often face inequities in resources, rest, and access to preventive care. Individuals from minoritized

communities or with low socioeconomic status often have significant non-paid caregiving roles that lead to inadequate sleep and poorer sleep quality.^{83,84} Similarly, minoritized communities often experience differences in neighborhood quality due to historical segregation, and resulting neighborhood disadvantage has been shown to partially account for race differences in sleep.⁸⁵ Poorer sleep is, in turn, associated with a higher overall inflammatory profile,^{86,87} and mediation analyses suggest that poorer global sleep quality underlies the link between major discrimination experience and inflammation burden. Inadequate sleep also leads to detrimental late-life cognitive outcomes due to poorer clearance of brain metabolites caused by altered functioning of the glymphatic system, a sleep-dependent pathway for interstitial fluid drainage.⁸⁸ In this context, our finding of greater age-related increases in interstitial water content among individuals exposed to major discrimination is noteworthy. Elevated interstitial water, as indexed by higher free water or isotropic volume fraction metrics from diffusion imaging, has been interpreted as a marker of altered interstitial fluid dynamics and is increasingly linked to neuroinflammation, cellular atrophy, impaired glymphatic clearance, and neurodegeneration.^{47,50,89} Although we did not directly examine sleep or glymphatic function in this project, these pathways offer a compelling hypothesis for how psychosocial adversity may contribute to brain aging and warrant further longitudinal and multimodal investigation.

Despite the considerable burden on those facing discrimination, it is important to note that individuals may have some agency in ameliorating its impact, and some individuals may grow after encountering discriminatory events and become more resilient to life's adversities.^{90–92} Building strong sources of social support,⁹³ being part of religious or spiritual communities,⁹⁴ engaging in active coping strategies such as values affirmation,⁹⁵ and practicing meditation,⁹⁶ may buffer against, at least some harmful effects of discrimination. These factors warrant future research in the context of brain microstructural changes in aging and potential resilience even in the face of adversity and discriminatory experiences.

Our study has several limitations that need to be considered. Firstly, our analyses are cross-sectional and do not permit causal inference. Secondly, consistent with existing literature such as in the Health and Retirement Study, there were few individuals with high levels of multiple major discriminatory experiences in our sample.⁷⁴ We therefore dichotomized discrimination experience into groups who reported experiencing at least one major discriminatory event compared with those who had no major experiences of discrimination, and thus we cannot infer dose-dependent effects of discrimination on brain health. Nevertheless, it is likely that the accelerated aging we observe in our sample may be considerably worse in a sample with higher levels of major discrimination across their lifetime. Thirdly, although diffusion measures from DTI, DKI, WMTI, and NODDI models have all been reported to be sensitive to age-related microstructural changes,^{20,62,97,98} they differ in their interpretability.¹⁹ Metrics derived from the statistical

models (DTI and DKI) have been widely reported to be sensitive to changes due to normal aging and pathologic changes, but as they can be influenced by multiple processes, they do not permit precise interpretation of the underlying biological changes,^{16,99} whereas biophysical models of diffusion such as NODDI and WMTI using multiple compartments provide more interpretable information about neuronal tissue microstructure.¹⁰⁰ Therefore, the biophysical models do not supersede the statistical ones, rather they refine and validate the interpretation. Each of these approaches make assumptions that need to be considered, none of them offer a comprehensive picture or are unequivocally better than the others, and newer improved models continue to be developed.^{53,101,102} Finally, we tested these diffusion models only in the whole-brain white matter skeleton and the left and right hippocampal regions, but not within the whole-brain gray matter. Future investigations of the association between discrimination and gray matter microstructure are warranted. Despite these limitations, the robust converging evidence we see from the DTI, NODDI, and WMTI diffusion MRI models, and lack of findings with DKI models, offers insights into the likely neurobiological pathways through which discrimination may be exerting adverse effects.

The next wave of MIDUS data collection in the same participants includes follow-up assessments of the measures included in this study as well as new measures of vascular imaging and both blood-based and positron emission tomography (PET) biomarkers for Alzheimer disease pathology. Similar data have been collected in the MIDUS Refresher cohort, which will further expand the study sample. The combined data set will provide sufficient power to examine interactions between discrimination, sex, education, and race. In addition to investigating longitudinal changes in DWI metrics, future investigations using these additional measures^{103,104} will clarify and help determine the mediating processes through which discrimination impacts brain health.

In summary, our findings suggest that experiencing at least one major discriminatory event is associated with accelerated brain microstructural aging. These data add to the accumulating evidence of the detrimental and widespread impacts of experiencing discrimination, ranging from the personal to the societal level. It is crucial to develop and disseminate effective tools and supportive interventions to reduce the impact of experiencing discrimination at the individual level. However, the burden of change should not solely rest on the individuals already coping with discrimination. Rather, given the high prevalence of discriminatory experiences, our findings underscore the need for policy interventions⁹⁵ to address and reduce discriminatory experiences that contribute to disparities in healthy and active aging, including brain health.

Acknowledgements: The authors are extremely grateful to the Midlife in the US (MIDUS) participants

and staff for their contributions to the study without which this research would not be possible.

Source of Funding and Conflicts of Interest: Publicly available data from the MIDUS study were used for this research. Since 1995, the MIDUS study has been funded by the following: the John D. and Catherine T. MacArthur Foundation Research Network, and the National Institute on Aging (P01-AG020166, U19-AG051426, U01-AG077928). The MIDUS Neuroscience Project's data collection was also supported in part by a core grant to the UW-Madison Waisman Center from the National Institute of Child Health and Human Development (P50HD105353). A.J.F. was supported by funding from the National Institute of Mental Health (F32-MH126537). R.J.D. is the founder, president, and serves on the board of directors for the non-profit organization, Healthy Minds Innovations, Inc. A.L.A. is part-owner and scientific advisor for ImgGyd, LLC which is developing technologies for image-guided surgeries. For the remaining authors none were declared.

Author Contributions: A.K.N.: writing—original draft, software, data curation, methodology, investigation, formal analysis, visualization. N.A.: writing—review and editing, software, data curation, methodology, investigation, visualization. A.J.F.: writing—review and editing, investigation. L.K.G.: writing—review and editing, data collection, data curation, project administration. S.E.S.: writing—review and editing, data collection, data curation. S.R.K.: writing—review and editing, methodology. A.L.A.: writing—review and editing, investigation. R.J.D.: writing—review and editing, funding acquisition. D.R.W.: writing—review and editing. C.D.R.: writing—review and editing, funding acquisition. S.M.S.: writing—review and editing, investigation, funding acquisition, conceptualization, resources, supervision.

Data Availability: All MIDUS data are archived and made publicly available—and are thus share-able—through the University of Michigan Inter-university Consortium of Political and Social Research (ICPSR), the MIDUS Portal (<https://midus.colectica.org/>), or through restricted access to the MIDUS Neuroscience repository (see https://www.midus.wisc.edu/midus_neuro_data.php/ for how to obtain access).

REFERENCES

1. Williams DR, Lawrence JA, Davis BA, Vu C. Understanding how discrimination can affect health. *Health Serv Res.* 2019;54:1374–1388.
2. Forde AT, Crookes DM, Suglia SF, Demmer RT. The weathering hypothesis as an explanation for racial disparities in health: a systematic review. *Ann Epidemiol.* 2019;33:1–18.e3.
3. Leger KA, Gloger EM, Maras J, Marshburn CK. Discrimination and health: the mediating role of daily stress processes. *Health Psychol.* 2022;41:332–342.
4. Brown TT, Partanen J, Chuong L, Villaverde V, Chantal Griffin A, Mendelson A. Discrimination hurts: the effect of discrimination on the development of chronic pain. *Soc Sci Med.* 2018;204:1–8.
5. Cuevas AG, Williams DR. Perceived Discrimination and Health: Integrative Findings. In: Ryff CD, Krueger RF. eds. *Oxf Handb Integr Health Sci.* Oxford University Press; 2018:442–456.
6. Whitman ET, Ryan CP, Abraham WC, Addae A, Corcoran DL,

- Elliott ML, et al. A blood biomarker of the pace of aging is associated with brain structure: replication across three cohorts. *Neurobiol Aging*. 2024;136:23–33.
7. Carroll JE, Price JE, Brown J, Bamishigbin O, Shalowitz MU, Ramey S, et al. Lifetime discrimination in low to middle income mothers and cellular aging: a prospective analysis. *Soc Sci Med*. 2022;311:115356.
 8. Cuevas AG, Cole SW, Belsky DW, McSorley AM, Shon JM, Chang VW. Multi-discrimination exposure and biological aging: Results from the midlife in the United States study. *Brain Behav Immun Health*. 2024;39:100774–100774.
 9. Zahodne LB, Sharifian N, Kraal AZ, Morris EP, Sol K, Zaheed AB, et al. Longitudinal associations between racial discrimination and hippocampal and white matter hyperintensity volumes among older Black adults. *Soc Sci Med*. 2023;316:114789.
 10. Barnes LL, Lewis TT, Begeny CT, Yu L, Bennett DA, Wilson RS. Perceived discrimination and cognition in older African Americans. *J Int Neuropsychol Soc*. 2012;18:856–865.
 11. Lindert J, Paul KC, Lachman Margie E, Ritz B, Seeman T. Social stress and risk of declining cognition: a longitudinal study of men and women in the United States. *Soc Psychiatry Psychiatr Epidemiol*. 2022;57:1875–1884.
 12. Bethlehem RAI, Seidlitz J, White SR, Vogel JW, Anderson KM, Adamson C, et al. Brain charts for the human lifespan. *Nature*. 2022;604:525–533.
 13. Fjell AM, Westlye LT, Grydeland H, Amlien I, Espeseth T, Reinvang I, et al. Critical ages in the life course of the adult brain: nonlinear subcortical aging. *Neurobiol Aging*. 2013;34:2239–2247.
 14. Merluzzi AP, Dean DC, Adluru N, Suryawanshi GS, Okonkwo OC, Oh JM, et al. Age-dependent differences in brain tissue microstructure assessed with neurite orientation dispersion and density imaging. *Neurobiol Aging*. 2016;43:79–88.
 15. Weston PSJ, Simpson IJA, Ryan NS, Ourselin S, Fox NC. Diffusion imaging changes in grey matter in Alzheimer's disease: a potential marker of early neurodegeneration. *Alzheimers Res Ther*. 2015;7:47.
 16. Alexander AL, Hurlley SA, Samsonov AA, Adluru N, Hosseinbor AP, Mossahebi P, et al. Characterization of cerebral white matter properties using quantitative magnetic resonance imaging stains. *Brain Connect*. 2011;1:423–446.
 17. Fieremans E, Jensen JH, Helpert JA. White matter characterization with diffusional kurtosis imaging. *NeuroImage*. 2011;58:177–188.
 18. Zhang H, Schneider T, Wheeler-Kingshott CA, Alexander DC. NODDI: practical in vivo neurite orientation dispersion and density imaging of the human brain. *NeuroImage*. 2012;61:1000–1016.
 19. DiPiero M, Rodrigues PG, Gromala A, Dean DC. Applications of advanced diffusion MRI in early brain development: a comprehensive review. *Brain Struct Funct*. 2022;228:367–392.
 20. Benítez A, Jensen JH, Thorn K, Dhiman S, Fountain-Zaragoza S, Rieter WJ, et al. Greater diffusion restriction in white matter in preclinical Alzheimer disease. *Ann Neurol*. 2022;91:864–877.
 21. Motovlyak A, Vogt NM, Adluru N, Ma Y, Wang R, Oh JM, et al. Age-related differences in white matter microstructure measured by advanced diffusion MRI in healthy older adults at risk for Alzheimer's disease. *Aging Brain*. 2022;2:100030.
 22. Nair AK, Adluru N, Finley AJ, Gresham LK, Skinner SE, Alexander AL, et al. Purpose in life as a resilience factor for brain health: diffusion MRI findings from the Midlife in the U.S. study. *Front Psychiatry*. 2024;15:1355998.
 23. Kessler RC, Mickelson KD, Williams DR. The prevalence, distribution, and mental health correlates of perceived discrimination in the United States. *J Health Soc Behav*. 1999;40:208.
 24. Williams DR, Yan Yu, Jackson JS, Anderson NB. Racial differences in physical and mental health: socio-economic status, stress and discrimination. *J Health Psychol*. 1997;2:335–351.
 25. Beatty Moody DL, Waldstein SR, Leibel DK, Hoggard LS, Gee GC, Ashe JJ, et al. Race and other sociodemographic categories are differentially linked to multiple dimensions of interpersonal-level discrimination: Implications for intersectional, health re-
- search. *PLOS ONE*. 2021;16:e0251174.
 26. Meza E, Peterson R, Gilsanz P, George KM, Miles SJ, Eng CW, et al. Perceived discrimination, nativity, and cognitive performance in a multiethnic study of older adults: findings from the Kaiser Healthy Aging and Diverse Life Experiences Study. *J Gerontol Ser A*. 2022;77:e65–e73.
 27. Williams DR, Gonzalez HM, Williams S, Mohammed SA, Moomal H, Stein DJ. Perceived discrimination, race and health in South Africa. *Soc Sci Med*. 2008;67:441–452.
 28. Ades-Aron B, Veraart J, Kochunov P, McGuire S, Sherman P, Kellner E, et al. Evaluation of the accuracy and precision of the diffusion parameter Estimation with Gibbs and Noise removal pipeline. *NeuroImage*. 2018;183:532–543.
 29. Jenkinson M, Beckmann CF, Behrens TEJ, Woolrich MW, Smith SM. FSL. *NeuroImage*. 2012;62:782–790.
 30. Avants BB, Tustison NJ, Song G, Cook PA, Klein A, Gee JC. A reproducible evaluation of ANTs similarity metric performance in brain image registration. *NeuroImage*. 2011;54:2033–2044.
 31. Tournier J-D, Smith R, Raffelt D, Tabbara R, Dhollander T, Pietsch M, et al. MRtrix3: A fast, flexible and open software framework for medical image processing and visualisation. *NeuroImage*. 2019;202:116137.
 32. Veraart J, Sijbers J, Sunaert S, Leemans A, Jeurissen B. Weighted linear least squares estimation of diffusion MRI parameters: strengths, limitations, and pitfalls. *NeuroImage*. 2013;81:335–346.
 33. Zhang Z, Vernekar D, Qian W, Kim M. Non-local means based Rician noise filtering for diffusion tensor and kurtosis imaging in human brain and spinal cord. *BMC Med Imaging*. 2021;21:16.
 34. Tabesh A, Jensen JH, Ardekani BA, Helpert JA. Estimation of tensors and tensor-derived measures in diffusional kurtosis imaging. *Magn Reson Med*. 2011;65:823–836.
 35. Perrone D, Aelterman J, Pižurica A, Jeurissen B, Philips W, Leemans A. The effect of Gibbs ringing artifacts on measures derived from diffusion MRI. *NeuroImage*. 2015;120:441–455.
 36. Lee JE, Chung MK, Lazar M, DuBray MB, Kim J, Bigler ED, et al. A study of diffusion tensor imaging by tissue-specific, smoothing-compensated voxel-based analysis. *NeuroImage*. 2009;44:870–883.
 37. Fick RHJ, Wassermann D, Deriche R. The Dmipy Toolbox: diffusion MRI multi-compartment modeling and microstructure recovery made easy. *Front Neuroinformatics*. 2019;13:64.
 38. Tustison NJ, Cook PA, Holbrook AJ, Johnson HJ, Muschelli J, Devenyi GA, et al. The ANTsX ecosystem for quantitative biological and medical imaging. *Sci Rep*. 2021;11:9068.
 39. Winkler AM, Webster MA, Brooks JC, Tracey I, Smith SM, Nichols TE. Non-parametric combination and related permutation tests for neuroimaging. *Hum Brain Mapp*. 2016;37:1486–1511.
 40. Smith SM, Jenkinson M, Woolrich MW, Beckmann CF, Behrens TEJ, Johansen-Berg H, et al. Advances in functional and structural MR image analysis and implementation as FSL. *NeuroImage*. 2004;23:S208–S219.
 41. Smith SM, Jenkinson M, Johansen-Berg H, Rueckert D, Nichols TE, Mackay CE, et al. Tract-based spatial statistics: voxelwise analysis of multi-subject diffusion data. *NeuroImage*. 2006;31:1487–1505.
 42. Frazier JA, Chiu S, Breeze JL, Makris N, Lange N, Kennedy DN, et al. Structural brain magnetic resonance imaging of limbic and thalamic volumes in pediatric bipolar disorder. *Am J Psychiatry*. 2005;162:1256–1265.
 43. R Core Team. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria. 2024. URL. <https://www.R-project.org/>
 44. Bancks MP, Byrd GS, Caban-Holt A, Fitzpatrick AL, Forrester SN, Hayden KM, et al. Self-reported experiences of discrimination and incident dementia. *Alzheimers Dement*. 2023;19:3119–3128.
 45. Lawrence JA, Kawachi I, White K, Bassett MT, Williams DR. Associations between multiple indicators of discrimination and allostatic load among middle-aged adults. *Soc Sci Med*. 2022;298:114866.
 46. Markon KE, Mann F, Freilich C, Cole S, Krueger RF. Associations between epigenetic age acceleration and longitudinal

- measures of psychosocioeconomic stress and status. *Soc Sci Med*. 2024;352:116990.
47. Raghavan S, Reid RI, Przybelski SA, Lesnick TG, Graff-Radford J, Schwarz CG, et al. Diffusion models reveal white matter microstructural changes with ageing, pathology and cognition. *Brain Commun*. 2021;3:fcab106.
 48. Pasternak O, Sochen N, Gur Y, Intrator N, Assaf Y. Free water elimination and mapping from diffusion MRI. *Magn Reson Med*. 2009;62:717–730.
 49. Wang Y, Wang Q, Haldar JP, Yeh F-C, Xie M, Sun P, et al. Quantification of increased cellularity during inflammatory demyelination. *Brain*. 2011;134:3590–3601.
 50. Chad JA, Pasternak O, Salat DH, Chen JJ. Re-examining age-related differences in white matter microstructure with free-water corrected diffusion tensor imaging. *Neurobiol Aging*. 2018;71:161–170.
 51. Baijot J, Van Laethem D, Denissen S, Costers L, Cambron M, D'Haeseleer M, et al. Radial diffusivity reflects general decline rather than specific cognitive deterioration in multiple sclerosis. *Sci Rep*. 2022;12:21771.
 52. Fieremans E, Benitez A, Jensen JH, Falangola MF, Tabesh A, Deardorff RL, et al. Novel white matter tract integrity metrics sensitive to Alzheimer disease progression. *Am J Neuroradiol*. 2013;34:2105–2112.
 53. Jelescu IO, De Skowronski A, Geffroy F, Palombo M, Novikov DS. Neurite Exchange Imaging (NEXI): a minimal model of diffusion in gray matter with inter-compartment water exchange. *NeuroImage*. 2022;256:119277.
 54. Karat BG, Köhler S, Khan AR. Diffusion MRI of the Hippocampus. *J Neurosci Off J Soc Neurosci*. 2024;44:e1705232024.
 55. Karat BG, DeKraker J, Hussain U, Köhler S, Khan AR. Mapping the macrostructure and microstructure of the in vivo human hippocampus using diffusion MRI. *Hum Brain Mapp*. 2023;44:5485–5503.
 56. Beaujoin J, Palomero-Gallagher N, Boumezbaur F, Axer M, Bernard J, Poupon F, et al. Post-mortem inference of the human hippocampal connectivity and microstructure using ultra-high field diffusion MRI at 11.7 T. *Brain Struct Funct*. 2018;223:2157–2179.
 57. Bettio LEB, Rajendran L, Gil-Mohapel J. The effects of aging in the hippocampus and cognitive decline. *Neurosci Biobehav Rev*. 2017;79:66–86.
 58. Terreros-Roncal J, Moreno-Jiménez EP, Flor-García M, Rodríguez-Moreno CB, Trinchero MF, Cafini F, et al. Impact of neurodegenerative diseases on human adult hippocampal neurogenesis. *Science*. 2021;374:1106–1113.
 59. Nazeri A, Schifani C, Anderson JAE, Ameis SH, Voineskos AN. In vivo imaging of gray matter microstructure in major psychiatric disorders: opportunities for clinical translation. *Biol Psychiatry Cogn Neurosci Neuroimaging*. 2020;5:855–864.
 60. Rahmanzadeh R, Lu P-J, Barakovic M, Weigel M, Maggi P, Nguyen TD, et al. Myelin and axon pathology in multiple sclerosis assessed by myelin water and multi-shell diffusion imaging. *Brain*. 2021;144:1684–1696.
 61. Assaf Y, Basser PJ. Composite hindered and restricted model of diffusion (CHARMED) MR imaging of the human brain. *NeuroImage*. 2005;27:48–58.
 62. Benitez A, Jensen JH, Falangola MF, Nietert PJ, Helpert JA. Modeling white matter tract integrity in aging with diffusional kurtosis imaging. *Neurobiol Aging*. 2018;70:265–275.
 63. Giampiccolo D, Herbet G, Duffau H. The inferior fronto-occipital fasciculus: bridging phylogeny, ontogeny and functional anatomy. *Brain*. 2025;148:1507–1525.
 64. Wang Y, Metoki A, Alm KH, Olson IR. White matter pathways and social cognition. *Neurosci Biobehav Rev*. 2018;90:350–370.
 65. Stoller F, Hinds E, Ionescu T, Khatamsaz E, Marston HM, Hengerer B. Forceps minor control of social behaviour. *Sci Rep*. 2024;14:30492.
 66. Ferris J, Greeley B, Yeganeh NM, Rinat S, Ramirez J, Black S, et al. Exploring biomarkers of processing speed and executive function: the role of the anterior thalamic radiations. *NeuroImage Clin*. 2022;36:103174.
 67. Potter LN, Brondolo E, Smyth JM. Biopsychosocial correlates of discrimination in daily life: A review. *Stigma Health*. 2019;4:38–61.
 68. Dark HE, Huang A, Cordon J, Deal JA, Palta P, Windham BG, et al. The association of perceived discrimination with dementia risk in Black older adults. *Alzheimers Dement*. 2023;19:4346–4356.
 69. Livingston G, Huntley J, Liu KY, Costafreda SG, Selbæk G, Alladi S, et al. Dementia prevention, intervention, and care: 2024 report of the Lancet standing Commission. *The Lancet*. 2024;404:572–628.
 70. Dong TS, Gee GC, Beltran-Sanchez H, Wang M, Osadchiy V, Kilpatrick LA, et al. How discrimination gets under the skin: biological determinants of discrimination associated with dysregulation of the brain-gut microbiome system and psychological symptoms. *Biol Psychiatry*. 2023;94:203–214.
 71. Oh H, Stickley A, Koyanagi A, Yau R, DeVlyder JE. Discrimination and suicidality among racial and ethnic minorities in the United States. *J Affect Disord*. 2019;245:517–523.
 72. Mann FD, Cuevas AG, Krueger RF. Cumulative stress: a general “s” factor in the structure of stress. *Soc Sci Med*. 2021;289:114405.
 73. Zahodne LB, Kraal AZ, Sharifian N, Zaheed AB, Sol K. Inflammatory mechanisms underlying the effects of everyday discrimination on age-related memory decline. *Brain Behav Immun*. 2019;75:149–154.
 74. Dhingra R, Hillmann AR, Reed RG. Major experiences of perceived discrimination across life and biological aging. *Psychoneuroendocrinology*. 2025;174:107380.
 75. Clark US, Miller ER, Hegde RR. Experiences of discrimination are associated with greater resting amygdala activity and functional connectivity. *Biol Psychiatry Cogn Neurosci Neuroimaging*. 2018;3:367–378.
 76. Sawyer PJ, Major B, Casad BJ, Townsend SSM, Mendes WB. Discrimination and the stress response: psychological and physiological consequences of anticipating prejudice in interethnic interactions. *Am J Public Health*. 2012;102:1020–1026.
 77. Meyer CS, Schreiner PJ, Lim K, Battapady H, Launer LJ. Depressive symptomatology, racial discrimination experience, and brain tissue volumes observed on magnetic resonance imaging. *Am J Epidemiol*. 2019;188:656–663.
 78. Sternthal MJ, Slopen N, Williams DR. Racial disparities in health: how much does stress really matter? *Bois Rev Soc Sci Res Race*. 2011;8:95–113.
 79. Callaghan BL, Tottenham N. The Stress Acceleration Hypothesis: effects of early-life adversity on emotion circuits and behavior. *Curr Opin Behav Sci*. 2016;7:76–81.
 80. Ellis BJ, Del Giudice M. Developmental adaptation to stress: an evolutionary perspective. *Annu Rev Psychol*. 2019;70:111–139.
 81. Hackman DA, Cserbik D, Chen J-C, Berhane K, Minaravesh B, McConnell R, et al. Association of local variation in neighborhood disadvantage in metropolitan areas with youth neurocognition and brain structure. *JAMA Pediatr*. 2021;175:e210426.
 82. McEwen BS, Gianaros PJ. Central role of the brain in stress and adaptation: Links to socioeconomic status, health, and disease. *Ann N Y Acad Sci*. 2010;1186:190–222.
 83. Gee GC, Hing A, Mohammed S, Tabor DC, Williams DR. Racism and the Life Course: Taking Time Seriously. *Am J Public Health*. 2019;109:S43–S47.
 84. Slopen N, Lewis TT, Williams DR. Discrimination and sleep: a systematic review. *Sleep Med*. 2016;18:88–95.
 85. Fuller-Rowell TE, Curtis DS, El-Sheikh M, Chae DH, Boylan JM, Ryff CD. Racial disparities in sleep: the role of neighborhood disadvantage. *Sleep Med*. 2016;27:281–8.
 86. Ong AD, Williams DR. Lifetime discrimination, global sleep quality, and inflammation burden in a multiethnic sample of middle-aged adults. *Cultur Divers Ethnic Minor Psychol*. 2019;25:82–90.
 87. Yip T, Chen M, Wang Y, Slopen N, Chae D, Priest N, et al. Linking discrimination and sleep with biomarker profiles: an investigation in the MIDUS study. *Compr Psychoneuroendocrinology*. 2021;5:100021.
 88. Nedergaard M, Goldman SA. Glymphatic failure as a final common pathway to dementia. *Science*. 2020;370:50–56.

89. Drenthen GS, Elschot EP, Van Der Knaap N, Uher D, Voorter PHM, Backes WH, et al. Imaging interstitial fluid with MRI: a narrative review on the associations of altered interstitial fluid with vascular and neurodegenerative abnormalities. *J Magn Reson Imaging*. 2024;60:40–53.
90. Grasser LR, Jovanovic T. Neural impacts of stigma, racism, and discrimination. *Biol Psychiatry Cogn Neurosci Neuroimaging*. 2022;7:1225–1234.
91. Ryff CD. The privilege of well-being in an increasingly unequal society. *Curr Dir Psychol Sci*. 2024;33:300–307.
92. Ryff CD, Keyes CLM, Hughes DL. Status inequalities, perceived discrimination, and eudaimonic well-being: do the challenges of minority life hone purpose and growth? *J Health Soc Behav*. 2003;44:275.
93. Hsieh N, Liu H, Zhang Z. Perceived discrimination and incident dementia among older adults in the United States: the buffering role of social relationships. *J Gerontol B Psychol Sci Soc Sci*. 2024;79:gbae059.
94. Ellison CG, Fan D. Daily spiritual experiences and psychological well-being among US adults. *Soc Indic Res*. 2008;88:247–271.
95. Lewis TT, Cogburn CD, Williams DR. Self-reported experiences of discrimination and health: scientific advances, ongoing controversies, and emerging issues. *Annu Rev Clin Psychol*. 2015;11:407–440.
96. Shallcross AJ, Spruill TM. The protective role of mindfulness in the relationship between perceived discrimination and depression. *Mindfulness*. 2018;9:1100–1109.
97. Lawrence KE, Nabulsi L, Santhalingam V, Abaryan Z, Villalón-Reina JE, Nir TM, et al. Age and sex effects on advanced white matter microstructure measures in 15,628 older adults: a UK biobank study. *Brain Imaging Behav*. 2021;15:2813–2823.
98. Schilling KG, Archer D, Yeh F-C, Rheault F, Cai LY, Hansen C, et al. Aging and white matter microstructure and macrostructure: a longitudinal multi-site diffusion MRI study of 1218 participants. *Brain Struct Funct*. 2022;227:2111–2125.
99. Jelescu IO, Zurek M, Winters KV, Veraart J, Rajaratnam A, Kim NS, et al. In vivo quantification of demyelination and recovery using compartment-specific diffusion MRI metrics validated by electron microscopy. *NeuroImage*. 2016;132:104–114.
100. Jelescu IO, Budde MD. Design and validation of diffusion MRI models of white matter. *Front Phys*. 2017;5:61.
101. Alsameen MH, Gong Z, Qian W, Kiely M, Triebswetter C, Bergeron CM, et al. C-NODDI: a constrained NODDI model for axonal density and orientation determinations in cerebral white matter. *Front Neurol*. 2023;14:1205426.
102. Palombo M, Ianus A, Guerreri M, Nunes D, Alexander DC, Shemesh N, et al. SANDI: a compartment-based model for non-invasive apparent soma and neurite imaging by diffusion MRI. *NeuroImage*. 2020;215:116835.
103. Kreisl WC, Kim M-J, Coughlin JM, Henter ID, Owen DR, Innis RB. PET imaging of neuroinflammation in neurological disorders. *Lancet Neurol*. 2020;19:940–950.
104. Miller KB, Howery AJ, Rivera-Rivera LA, Johnson SC, Rowley HA, Wieben O, et al. Age-related reductions in cerebrovascular reactivity using 4D flow MRI. *Front Aging Neurosci*. 2019;11:281.